

Continuous Self-Similar Evaporation of a Rotating Cosmic String Loop

Malcolm Anderson
 Department of Mathematics
 Universiti Brunei Darussalam
 Jalan Tungku Link, Gadong BE 1410
 Negara Brunei Darussalam

Abstract

A solution of the linearized Einstein and Nambu-Goto equations is constructed which describes the evaporation of a certain type of rotating cosmic string – the Allen-Casper-Ottewill loop – under the action of its own self-gravity. The solution evaporates self-similarly, and radiates away all its mass-energy and momentum in a finite time. Furthermore, the corresponding weak-field metric can be matched to a remnant Minkowski spacetime at all points on the future light cone of the final evaporation point of the loop.

Short Title: Evaporation of a String Loop

PACS numbers: 04.25.Nx, 98.80.Cq

1. Introduction

Cosmic strings are thin filaments of topologically-trapped Higgs field energy whose dynamics, in the zero-thickness limit, are controlled by the Nambu-Goto action. In this limit cosmic strings are effectively line singularities which move under the action of their own elastic tension, and radiate gravitational energy and momentum in the process. They may have played a role in the formation of large-scale structure in the early Universe (see [1] for a review), but their gravitational properties are still only poorly understood, and they consequently remain problematic

objects for general relativity. In particular, a rigorous distributional description of line singularities has not yet been developed within the framework of general relativity, nor has it been demonstrated that zero-thickness cosmic strings can always be interpreted as the unique limits of well-behaved finite-thickness solutions [2].

At present, all known exact self-gravitating string solutions in a Minkowski background describe either infinite straight strings [3, 4, 5, 6] or infinite strings interacting with plane-fronted gravitational waves [7, 8], and the only general result available on the dynamics of string *loops* in strong-field gravity is Hawking's proof that a collapsing circular loop would form an event horizon after radiating away at most 29% of its original energy [9]. (In addition, the full 3-metric outside a circular loop at a moment of time symmetry has been constructed in [10].)

Given the complexity of the general strong-field problem, therefore, the most promising arena for the study of loop self-gravity would appear to be the weak-field limit. Fortunately, it can be shown that the self-gravity of a GUT string would almost everywhere be small enough to justify a weak-field treatment [11], although this treatment typically breaks down in the vicinity of certain common pathological features known as kinks and cusps.

In [12] (henceforth referred to as Paper I) it was shown that a certain type of rigidly-rotating cosmic string loop – the Allen-Casper-Ottewill solution [13] – evolves by self-similar shrinkage in the weak-field limit. This is the first explicit back-reaction calculation to have been published for a cosmic string loop, as all previous calculations have relied on numerical approximation [11]. However, the calculation relies on what might be called an adiabatic approximation, as the total flux of gravitational energy and momentum from the loop is integrated over a single oscillation period of the loop's unperturbed trajectory, and is shown to induce no changes other than a small decrease in the loop's energy and length, plus a comparable rotational phase shift.

In this paper I extend the analysis of Paper I by relaxing the adiabatic assumption and constructing a self-similar evaporating solution of the linearized field equations which is identical to the Allen-Casper-Ottewill solution on spacelike sections and ultimately radiates away all its mass-energy. Although the trajectory of the string loop, and the perturbations it induces in the background metric, are only minimally different from the corresponding quantities in the adiabatic case, the evaporating solution represents an incremental advance in the understanding of loop self-gravity. In particular, the metric perturbations can be calculated at all points outside the string's worldsheet, and the perturbed (radiating) spacetime

can be matched to a flat spacetime which occupies the causal future of the final point of evaporation of the loop. The naive expectation that the loop evaporates completely to leave behind a vacuum remnant spacetime therefore appears to be consistent at this level of approximation.

2. Preliminaries

The mathematical formalism used to describe zero-thickness cosmic strings has been discussed in some detail in Paper I, and in this section I will offer only a brief summary.

Throughout the paper, $x^a \equiv [x^0, x^1, x^2, x^3] = [t, x, y, z]^a$ are local coordinates on the four-dimensional background spacetime $(\mathbf{M}, g_{ab})$, and the metric tensor g_{ab} has signature $(+, -, -, -)$, so that timelike vectors have positive norm. The world sheet $\mathbf{T}$ of the string is the two-dimensional surface it traces out as it moves, and is described parametrically by a set of equations of the form $x^a = X^a(\zeta^A)$, where the parameters $\zeta^A \equiv (\zeta^0, \zeta^1)$ are gauge coordinates.

The intrinsic two-metric induced on $\mathbf{T}$ by a given choice of gauge coordinates is:

$$\gamma_{AB} = g_{ab} X^a{}_{,A} X^b{}_{,B} \quad (2.1)$$

where $X^a{}_{,A}$ is shorthand for $\partial X^a / \partial \zeta^A$. The intrinsic metric γ_{AB} is assumed to be non-degenerate with signature $(+, -)$ almost everywhere on $\mathbf{T}$. If γ denotes $|\det(\gamma_{AB})|$ the Nambu-Goto action [14, 15] has the form

$$I = -\mu \int \gamma^{1/2} d^2\zeta \quad (2.2)$$

where μ is the (constant) mass per unit length of the string.

The stress-energy tensor T^{ab} induced by a given trajectory is found by varying the Nambu-Goto action with respect to the background metric g_{ab} , according to the standard prescription $\delta I = -\frac{1}{2} \int T^{ab} \delta g_{ab} g^{1/2} d^4x$, where $g \equiv |\det(g_{ab})|$. The equation for δI inverts to give

$$T^{ab}(x^c) = -2g^{-1/2} \int (\delta \mathcal{L} / \delta g_{ab}) \delta^{(4)}(x^c - X^c) d^2\zeta \quad (2.3)$$

where $\mathcal{L} \equiv -\mu \gamma^{1/2}$ is the Lagrangian density in (2.2), and

$$\delta \mathcal{L} / \delta g_{ab} = -\frac{1}{2} \mu \gamma^{1/2} \gamma^{AB} X^a{}_{,A} X^b{}_{,B} \quad (2.4)$$

with γ^{AB} the inverse of γ_{AB} .

In Paper I the gauge choice $\gamma_{AB} = \kappa\eta_{AB}$ was imposed from the outset, where $\eta_{AB} = \text{diag}(1, -1)$ is the Minkowski 2-tensor. However, it turns out that this choice is not as convenient when the string is evaporating (and the trajectory is no longer time-periodic), and for this reason the more general versions of the string stress-energy and equation of motion will be retained in what follows.

The extrinsic curvature tensor of the world sheet $\mathbf{T}$ has the general form

$$K^{abc} = p_m^a p^{bn} \nabla_n p^{mc} \quad (2.5)$$

where ∇_n is the derivative operator associated with g_{ab} , and $p^{ab} \equiv \gamma^{AB} X^a_{,A} X^b_{,B}$ is the projection tensor corresponding to γ_{AB} . In terms of the derivatives of the position function X^a the curvature tensor K^{abc} and its trace $K^c \equiv g_{ab} K^{abc}$ can be written as

$$K^{abc} = q_d^c (\gamma^{AC} \gamma^{BD} X^a_{,A} X^b_{,B} X^d_{,CD} + p^{am} p^{bn} \Gamma_{mn}^d) \quad (2.6)$$

and

$$K^c = q_d^c (\gamma^{CD} X^d_{,CD} + p^{mn} \Gamma_{mn}^d) \quad (2.7)$$

with $q^{ab} \equiv g^{ab} - p^{ab}$ the orthogonal complement of p^{ab} , and Γ_{bc}^a the Christoffel symbol associated with g_{ab} . The equation of motion of the string is recovered by setting the functional derivative $\delta\mathcal{L}/\delta X^a$ equal to zero, and reads simply:

$$K^c = 0 \quad (2.8)$$

In the absence of any sources of stress-energy other than the string itself, the system of equations for the metric and the string's trajectory is closed by imposing the Einstein equation $G^{ab} = -8\pi T^{ab}$, where $G^{ab} = R^{ab} - \frac{1}{2}g^{ab}R$ is the Einstein tensor.¹ Equations (2.3), (2.4) and (2.8), together with the Einstein equation, constitute the strong-field back-reaction problem. As was mentioned above, the only known solutions to the strong-field back-reaction problem describe either infinite straight strings (with or without an envelope of cylindrical gravitational waves [3, 4, 5, 6]) or infinite strings interacting with plane-fronted gravitational waves (the so-called travelling-wave solutions [7, 8]). There are no known exact loop

¹The convention adopted here for the Riemann tensor is that

$$R_{abcd} = \frac{1}{2}(g_{ac,bd} + g_{bd,ac} - g_{ad,bc} - g_{bc,ad}) + g_{ef}(\Gamma_{ac}^e \Gamma_{bd}^f - \Gamma_{ad}^e \Gamma_{bc}^f).$$

Also, geometrized units have been chosen, so that the gravitational constant G and the speed of light c are set to 1.

solutions in a vacuum background, and an analysis of loop evolution is currently feasible only at the level of the weak-field approximation.

In the weak-field formalism the field and trajectory variables are truncated at linear order in the mass per unit length μ of the string, which is assumed to be small compared to 1. This is certainly thought to be true of GUT strings, for which $\mu \sim 10^{-6}$. The underlying assumption is that the solution pair (g_{ab}, X^a) to the strong-field back-reaction problem can be expanded as perturbative series of the form

$$g_{ab} = \sum_{k=0}^{\infty} \mu^k g_{ab}^{(k)} \quad \text{and} \quad X^a = \sum_{k=0}^{\infty} \mu^k X_{(k)}^a \quad (2.9)$$

where the functions $g_{ab}^{(k)}$ and $X_{(k)}^a$ are independent of μ . The weak-field back-reaction problem consists in setting $g_{ab}^{(0)}$ equal to η_{ab} , the Minkowski metric tensor, and solving the Einstein equation and equation of motion (2.8) at linear order in μ to obtain the metric perturbation $g_{ab}^{(1)}$ and the trajectory functions $X_{(0)}^a$ and $X_{(1)}^a$.

In what follows, the Minkowski 4-tensor will be taken to have its rigid form $\eta_{ab} = \text{diag}(1, -1, -1, -1)$. In solving the weak-field problem, a gauge must first be chosen for the metric perturbation $g_{ab}^{(1)}$, and also for the functions $X_{(0)}^a$ and $X_{(1)}^a$. If $h_{ab} \equiv \mu g_{ab}^{(1)}$ the standard harmonic gauge conditions read $h_{a,b}^b = \frac{1}{2} h_{,a}$ (where $h \equiv h_b^b$ and indices are everywhere lowered and raised using η_{ab} and its inverse η^{ab}). The Einstein equation $G^{ab} = -8\pi T^{ab}$ then reduces to $\square h_{ab} = -16\pi S_{ab}$, where $S_{ab} = T_{ab} - \frac{1}{2}\eta_{ab}T_c^c$ and $\square \equiv \partial_t^2 - \nabla^2$ is the flat-space d'Alembertian.

With $x^a \equiv [t, \mathbf{x}]^a$ the retarded solution for h_{ab} is:

$$h_{ab}(t, \mathbf{x}) = -4 \int \frac{S_{ab}(t', \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} d^3x' \quad (2.10)$$

where $t' = t - |\mathbf{x} - \mathbf{x}'|$ is the retarded time at the source point $\mathbf{x}'$. Since $g^{1/2} = 1$ to leading order in h_{ab} , equations (2.3) and (2.4) together give

$$S_{ab}(x'^d) = \mu \int \gamma^{1/2} \gamma^{AB} (X_{(0)a,A} X_{(0)b,B} - \frac{1}{2} \eta_{ab} X_{(0),A}^c X_{(0)c,B}) \delta^{(4)}(x'^d - X_{(0)}^d) d^2\zeta \quad (2.11)$$

with $\gamma^{1/2}$ and γ^{AB} evaluated using η_{ab} in place of g_{ab} and $X_{(0)}^a$ in place of X^a .

Furthermore, if $g_{ab}^{(0)} = \eta_{ab}$ then $\Gamma_{mn}^d = 0$ to leading order, and at linear order

in h_{ab} the equation of motion $K^c = 0$ reads:

$$q_d^c (\gamma^{CD} X^d_{,CD} - \gamma^{AC} \gamma^{BD} h_{ab} X^a_{,A} X^b_{,B} X^d_{,CD}) = -q^{cd} \gamma^{AB} X^a_{,A} X^b_{,B} (h_{bd,a} - \frac{1}{2} h_{ab,d}) \quad (2.12)$$

where q_d^c and γ^{AC} are evaluated using η_{ab} rather than g_{ab} (but X^a is retained in full). As was explained in Paper I, this equation is just the flat-space version of the Battye-Carter equation [16, 17].

If X^a is now replaced by $X^a_{(0)} + \delta X^a$ where $\delta X^a \equiv \mu X^a_{(1)}$ then, with q_d^c and γ^{AB} evaluated using $X^a_{(0)}$ in place of X^a , equation (2.12) splits into two parts:

$$q_d^c \gamma^{CD} X^d_{(0),CD} = 0 \quad (2.13)$$

and

$$\begin{aligned} & q_d^c \gamma^{CD} \delta X^d_{,CD} - 2\eta_{ab} \gamma^{AB} q_a^{(b} X^c_{(0),A} \delta X^a_{,B} \gamma^{CD} X^d_{(0),CD} \\ & - q_d^c \gamma^{AC} \gamma^{BD} (2\eta_{ab} X^a_{(0),A} \delta X^b_{,B} + h_{ab} X^a_{(0),A} X^b_{(0),B}) X^d_{(0),CD} \\ = & -q^{cd} \gamma^{AB} X^a_{(0),A} X^b_{(0),B} (h_{bd,a} - \frac{1}{2} h_{ab,d}) \end{aligned} \quad (2.14)$$

at zeroth order and first order in μ respectively. (Here, round brackets on indices denote symmetrization, so that $q_a^{(b} X^c_{(0),A} \equiv \frac{1}{2} [q_a^b X^c_{(0),A} + q_a^c X^b_{(0),A}]$.)

The weak-field back-reaction problem, in its most general form, therefore reduces to equations (2.10), (2.13) and (2.14) for h_{ab} , $X^a_{(0)}$ and δX^a . In order to specialize these equations to the problem in hand, it is necessary to first explain the choice that will be made for $X^a_{(0)}$, and the corresponding choice of gauge coordinates ζ^0 and ζ^1 . This will be done in Sections 3 and 4 below.

3. The Allen-Casper-Ottewill Loop

The equation of motion at leading order (2.13) can be solved exactly by making a suitable choice of gauge coordinates. A common choice is the so-called standard gauge $\zeta^A = [\tau, \sigma]^A$, in which γ_{AB} is everywhere proportional to the Minkowski tensor $\eta_{AB} = \text{diag}(1, -1)$. Since $\gamma_{AB} = \eta_{ab} X^a_{(0),A} X^b_{(0),B}$ the geometric significance of this gauge choice is that the tangent vectors $X^a_{(0),\tau}$ (which is timelike) and $X^a_{(0),\sigma}$ (which is spacelike) are orthogonal and of equal magnitude. As was demonstrated in Paper I, the equation of motion (2.13) reduces in the standard gauge to the wave equation

$$X^a_{(0),\tau\tau} - X^a_{(0),\sigma\sigma} = 0 \quad (3.1)$$

If the subsidiary gauge alignment condition $X_{(0)}^0(\tau, \sigma) = \tau$ is imposed then all loop solutions of (2.13) can be written in the form $X_{(0)}^a = [\tau, \mathbf{X}(\tau, \sigma)]^a$, where

$$\mathbf{X}(\tau, \sigma) = \frac{1}{2}[\mathbf{a}(\tau + \sigma) + \mathbf{b}(\tau - \sigma)] \quad (3.2)$$

for some choice of vector functions $\mathbf{a}$ and $\mathbf{b}$. The gauge constraints $\gamma_{\tau\tau} + \gamma_{\sigma\sigma} = 0$ and $\gamma_{\sigma\tau} = 0$ together imply that $|\mathbf{a}'|^2 = |\mathbf{b}'|^2 = 1$, where a prime denotes differentiation with respect to the relevant argument. If the loop is in its centre-of-momentum frame then $\mathbf{a}$ and $\mathbf{b}$ are each periodic functions of their arguments with some parametric period L , and since $\mathbf{X}(\tau + L/2, \sigma + L/2) = \mathbf{X}(\tau, \sigma)$ for any values of τ and σ when $\mathbf{a}$ and $\mathbf{b}$ are periodic, the fundamental oscillation period t_p of the loop is $L/2$.

In the centre-of-momentum frame, the total four-momentum of the string loop on any surface of constant $t = \tau$ is:

$$p^a = \mu \int_0^L X_{(0),\tau}^a d\sigma = \mu \int_0^L [1, \frac{1}{2}\mathbf{a}'(\tau + \sigma) + \frac{1}{2}\mathbf{b}'(\tau - \sigma)] d\sigma = \mu L[1, \mathbf{0}] \quad (3.3)$$

and in particular the energy of the loop is $E = \mu L$. The corresponding angular momentum of the loop is:

$$\mathbf{J} = \frac{1}{4}\mu \int_0^L (\mathbf{a} \times \mathbf{a}' + \mathbf{b} \times \mathbf{b}') d\sigma. \quad (3.4)$$

In the weak-field limit, the gravitational power P radiated by an oscillating compact source can be calculated from the standard expression for the power per unit solid angle in the wave zone (at distances large compared to the characteristic size L of the source):

$$\frac{dP}{d\Omega} = \frac{\omega^2}{\pi} \sum_{m=1}^{\infty} m^2 [\bar{T}^{ab} \bar{T}_{ab}^* - \frac{1}{2} |\bar{T}_a^a|^2] \quad (3.5)$$

where $\omega = 2\pi/t_p$ is the circular frequency of the source, and

$$\bar{T}^{ab}(m, \mathbf{n}) = t_p^{-1} \int_{\mathbb{R}^3} \int_0^{t_p} T^{ab}(t, \mathbf{x}) e^{im\omega(t - \mathbf{n} \cdot \mathbf{x})} dt d^3x \quad (3.6)$$

is the Fourier transform of its integrated stress-energy, with $\mathbf{n}$ the unit vector in the direction of the field point.

For a string loop in a flat background in the standard gauge, the stress-energy tensor (2.3) becomes

$$T^{ab}(x^c) = \mu \int (X_{(0),\tau}^a X_{(0),\tau}^b - X_{(0),\sigma}^a X_{(0),\sigma}^b) \delta^{(4)}(x^c - X_{(0)}^c) d\tau d\sigma \quad (3.7)$$

and the power per unit solid angle reduces to

$$\frac{dP}{d\Omega} = 8\pi\mu^2 \sum_{m=1}^{\infty} m^2 [(A \cdot A^*)(B \cdot B^*) + |A \cdot B^*|^2 - |A \cdot B|^2]. \quad (3.8)$$

where

$$A^c(m, \mathbf{n}) = \frac{1}{L} \int_0^L e^{2\pi i m [\sigma_+ - \mathbf{n} \cdot \mathbf{a}(\sigma_+)]/L} [1, \mathbf{a}'(\sigma_+)]^c d\sigma_+ \quad (3.9)$$

and

$$B^c(m, \mathbf{n}) = \frac{1}{L} \int_0^L e^{2\pi i m [\sigma_- - \mathbf{n} \cdot \mathbf{b}(\sigma_-)]/L} [1, \mathbf{b}'(\sigma_-)]^c d\sigma_-. \quad (3.10)$$

with the gauge coordinates σ_+ and σ_- defined by $\sigma_{\pm} \equiv \tau \pm \sigma$. (Here and elsewhere the dot product is taken with respect to the flat-space metric η_{ab} .)

The radiative efficiency γ^0 of a string loop is defined to be $\gamma^0 \equiv \mu^{-2} P$, where $P = \int \frac{dP}{d\Omega} d\Omega$ is the total power of the loop. The radiative efficiency has been calculated analytically for a large class of simple loop trajectories by Allen, Casper and Ottewill [13], and numerically for the loop by-products of string network simulations by Allen and Shellard [18] and Allen and Casper [19]. The simulation studies found that the value of γ^0 is typically of order 65-70, and seems to be bounded below by about 40. This observation is consistent with the results of the analytic study, which found that γ^0 has a minimum value of about 39.0025, and occurs if the mode functions have the forms

$$\mathbf{a}(\sigma_+) = (|\sigma_+| - \frac{1}{4}L) \hat{\mathbf{z}} \quad \text{for} \quad -\frac{1}{2}L \leq \sigma_+ \leq \frac{1}{2}L \quad (3.11)$$

and

$$\mathbf{b}(\sigma_-) = \frac{L}{2\pi} [\cos(2\pi\sigma_-/L) \hat{\mathbf{x}} + \sin(2\pi\sigma_-/L) \hat{\mathbf{y}}], \quad (3.12)$$

with $\mathbf{a}$ represented by its even periodic extension when σ_+ lies outside $[-\frac{1}{2}L, \frac{1}{2}L]$.²

²Strictly speaking, the $\mathbf{a}$ mode function was represented in both [13] and Paper I in the form

$$\mathbf{a}(\sigma_+) = \begin{cases} \sigma_+ \hat{\mathbf{z}} & \text{if } 0 \leq \sigma_+ < \frac{1}{2}L \\ (L - \sigma_+) \hat{\mathbf{z}} & \text{if } \frac{1}{2}L \leq \sigma_+ < L \end{cases}$$

The corresponding loop, which I will henceforth refer to as the Allen-Casper-Ottewill or ACO loop, is rigidly rotating about the z -axis with an angular speed $\omega = 4\pi/L$. The evolution of the loop is illustrated in Figure 1, which shows the y - z projection of the loop at times $\tau - \varepsilon = 0, L/16, L/8$ and $3L/16$ (top row) and $\tau - \varepsilon = L/4, 5L/16, 3L/8$ and $7L/16$ (bottom row), where the time offset ε is $0.02L$. The string has been artificially thickened for the sake of visibility, and the z -axis is also shown. The projections of the loop onto the x - y plane are circles of radius $L/(4\pi)$. The points at the extreme top and bottom of the loop, where the helical segments meet and the modal tangent vector $\mathbf{a}'$ is discontinuous, are technically known as kinks. They trace out circles in the planes $z = L/8$ and $z = -L/8$. All other points on the loop trace out identical circles, although with varying phase lags. The net angular momentum of the loop is $\mathbf{J} = \frac{1}{8\pi}\mu L^2 \hat{\mathbf{z}}$.

I should stress that it is not currently known whether the ACO solution has the lowest radiative efficiency of all possible string loops. Allen, Casper and Ottewill [13] showed only that the ACO solution minimizes γ^0 over a special class of solutions with the mode function $\mathbf{a}$ given by (3.11) and the mode function $\mathbf{b}$ confined to the x - y plane but otherwise arbitrary. Nonetheless, in Paper I the first-order equation of motion (2.14) was solved for a leading-order trajectory $X_{(0)}^a$ of ACO form, and it was shown that the net perturbation δX^a induced in the string trajectory over a single oscillation period $\Delta\tau = t_p$ corresponds to no more than a change $\Delta L = -\frac{1}{2}\gamma^0\mu L \approx -19.501\mu L$ in the length and parametric period L of the loop, and a rotational phase shift with magnitude about 38.92μ acting to advance the overall pattern of the loop. In other words, as it radiates the ACO loop retains its shape and therefore its original radiative efficiency. If the ACO loop does indeed possess the lowest possible radiative efficiency, it follows that it would be the longest-lived loop in any ensemble of loops with the same energy.

4. The Trajectory of the Evaporating ACO Loop

Whether or not the ACO loop minimizes γ^0 , the fact that γ^0 remains constant as the loop evaporates is the key to a more extended description of the loop's evolution. Over the course of a double period $\Delta t = 2t_p = L$ the total energy $E = \mu L$ of the loop is reduced by an amount $\Delta E = -PL = -\mu^2\gamma^0 L$ and so

in which case the centroid of the loop lies at $(x, y, z) = (0, 0, \frac{1}{8}L)$. For the purposes of the analysis presented below it is more convenient to translate the loop down the z -axis by an amount $\frac{1}{8}L$ so that the centroid lies at the origin. The resulting mode function is then given by (3.11).

$\Delta E/\Delta t = -\mu^2\gamma^0$, or equivalently $\Delta L/\Delta t = -\mu\gamma^0$. Since the loop is rotating uniformly, this relation is also true instantaneously, so that $dL/dt = -\mu\gamma^0$ and therefore $L(t) = L_0 - \mu\gamma^0(t - t_0)$, where L_0 is the value of L at some initial time t_0 . In particular, if the time coordinate is chosen so that $t_0 = 0$ then

$$L(t) = L_0(1 - t/t_L) \quad (4.1)$$

for $t < t_L$, where $t_L \equiv L_0/(\gamma^0\mu)$ is the total future lifetime of the loop.

If the adiabatic calculation performed in Paper I accurately represents the evolution of the ACO loop, it is expected that as it evaporates the loop retains the same spatial cross-section, apart from a uniform reduction in scale proportional to $L(t)$, and a rigid rotation about the z -axis. What follows now is a heuristic argument leading to a plausible form for the trajectory of the evaporating loop. It should be cautioned that the argument below relies implicitly on an assumption that the background spacetime is flat, and consideration of the first-order equation of motion (2.14) quickly shows that this assumption is overly optimistic. Nonetheless, it is also clear on physical grounds what form the relativistic corrections to the trajectory should have, and the first-order equation of motion will be solved to give these corrections explicitly in Section 6.

The most general form of the position function $\mathbf{X}$ of the evaporating loop (with τ henceforth replaced by the time coordinate t , and ξ a general parameter) is

$$\mathbf{X}(t, \xi) = L(t) \bar{\mathbf{X}}(t, \xi) \quad (4.2)$$

where $\bar{\mathbf{X}}$ is rigidly rotating, so that if $t < t_L$ and $t^* < t_L$ are two different times then

$$\bar{\mathbf{X}}(t, \xi) = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix} \bar{\mathbf{X}}(t^*, \xi^*) \quad (4.3)$$

with $\phi(t^*, t)$ a phase shift and $\xi^*(t, t^*, \xi)$ a reparametrization of ξ .

Here, if t^* is fixed, $\bar{\mathbf{X}}(t^*, \xi^*)$ is a reference function that at each value of t must trace out the same loop in $\mathbb{R}^3$ as ξ varies over its range. In particular, if $t \rightarrow t + \delta t$ then $\xi^*(t + \delta t, t^*, \xi + \delta \xi) = \xi^*(t, t^*, \xi)$ for some function $\delta \xi(\xi)$. That is,

$$\partial_t \xi^* \delta t + \partial_\xi \xi^* \delta \xi = 0 \quad (4.4)$$

to linear order in δt and $\delta \xi$, and since this is a homogeneous linear first-order PDE with characteristic $\xi = \alpha(t)$ for some function α , it follows that ξ^* can be

any function of $\xi - \alpha(t)$. Hence, if the fixed reference time t^* is suppressed,

$$\mathbf{X}(t, \xi) = L(t) \begin{bmatrix} \cos \phi(t) & -\sin \phi(t) & 0 \\ \sin \phi(t) & \cos \phi(t) & 0 \\ 0 & 0 & 1 \end{bmatrix} \bar{\mathbf{X}}_0(\xi - \alpha(t)) \quad (4.5)$$

for some choice of functions ϕ , $\bar{\mathbf{X}}_0$ and α .

Furthermore, because the parametric period of a cosmic string loop scales with its length L , all timescales on the loop do likewise, and in particular the velocity field $\mathbf{X}_t$ of the loop must at all times be rotating rigidly. Now,

$$\mathbf{X}_t = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix} [L' \bar{\mathbf{X}}_0 + L(\phi' M \bar{\mathbf{X}}_0 - \alpha' \bar{\mathbf{X}}_0')] \quad (4.6)$$

where

$$M = \begin{bmatrix} \cos \phi & \sin \phi & 0 \\ -\sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -\sin \phi & -\cos \phi & 0 \\ \cos \phi & -\sin \phi & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad (4.7)$$

and the primes denote derivatives with respect to the relevant arguments.

If the vector functions $\bar{\mathbf{X}}_0$, $M \bar{\mathbf{X}}_0$ and $\bar{\mathbf{X}}_0'$ are linearly independent, the requirement that $\mathbf{X}_t$ be rigidly rotating constrains L' , $L\phi'$ and $L\alpha'$ to all be constant. The first condition, that L' be constant, was derived independently at the beginning of this section, and is equivalent to requiring the radiative efficiency γ^0 to be constant. Since $L(t) = L_0(1 - t/t_L)$ for $t < t_L$, the remaining constraints can be integrated immediately to give

$$\phi(t) = -p \ln(1 - t/t_L) + \phi_0 \quad \text{and} \quad \alpha(t) = -q \ln(1 - t/t_L) + \alpha_0 \quad (4.8)$$

where p , q , ϕ_0 and α_0 are constants.

By suitably rotating the x - y coordinate axes and rezeroing the gauge coordinate ξ it is always possible to set ϕ_0 and α_0 to zero. Furthermore, if $\bar{\sigma} \equiv p\xi/q$ then $\bar{\mathbf{X}}_0$ is a function of $\bar{\sigma} + p \ln(1 - t/t_L)$ and so

$$\mathbf{X}(t, \bar{\sigma}) = L_0(1 - t/t_L) \begin{bmatrix} \cos \phi(t) & -\sin \phi(t) & 0 \\ \sin \phi(t) & \cos \phi(t) & 0 \\ 0 & 0 & 1 \end{bmatrix} \bar{\mathbf{X}}_0(\bar{\sigma} - \phi(t)) \quad (4.9)$$

with $\phi(t) = -p \ln(1 - t/t_L)$.

If this trajectory is to describe the evaporation of an ACO loop, the position function (4.9) and its time derivative should, to leading order in μ , match the position function $\mathbf{X}$ and velocity $\mathbf{X}_t$ of the unperturbed ACO loop at time $t = 0$. The ACO loop is described by the mode functions (3.11) and (3.12), so its position function has the explicit form

$$\mathbf{X}(t, \sigma) = \frac{1}{4\pi}L\{\cos[2\pi(t - \sigma)/L]\hat{\mathbf{x}} + \sin[2\pi(t - \sigma)/L]\hat{\mathbf{y}}\} + \frac{1}{2}(|t + \sigma| - \frac{1}{4}L)\hat{\mathbf{z}} \quad (4.10)$$

for $-\frac{1}{2}L \leq t + \sigma \leq \frac{1}{2}L$.

Here, the constant scale factors L_0 and L are taken to be identical. Since $\phi(0) = 0$, the position functions (4.9) and (4.10) will agree at $t = 0$ if

$$\bar{\mathbf{X}}_0(\bar{\sigma}) = \frac{1}{4\pi}L[\cos(2\pi\sigma/L)\hat{\mathbf{x}} - \sin(2\pi\sigma/L)\hat{\mathbf{y}}] + \frac{1}{2}(|\sigma| - \frac{1}{4}L)\hat{\mathbf{z}}. \quad (4.11)$$

for $-\frac{1}{2}L \leq \sigma \leq \frac{1}{2}L$. Furthermore, $L'(t) = -L_0/t_L$ is of order μ and can be ignored when calculating $\mathbf{X}_t$, while $\phi'(t) = p/t_L$ at $t = 0$. Equating the time derivatives of (4.9) and (4.10) at $t = 0$ then leads to the consistency conditions $\bar{\sigma} = -4\pi\sigma/L$ and $p = 4\pi t_L/L_0 \equiv 4\pi/(\gamma^0\mu)$. After substituting these conditions and equation (4.11) into (4.9), the position function of the evaporating loop becomes:

$$\mathbf{X}(t, \bar{\sigma}) = \frac{1}{4\pi}L_0(1 - t/t_L)\{\cos[\frac{1}{2}(\phi(t) + \bar{\sigma})]\hat{\mathbf{x}} + \sin[\frac{1}{2}(\phi(t) + \bar{\sigma})]\hat{\mathbf{y}} + \frac{1}{2}[|\phi(t) - \bar{\sigma}| - \pi]\hat{\mathbf{z}}\} \quad (4.12)$$

for $-2\pi \leq \phi(t) - \bar{\sigma} \leq 2\pi$, where $\phi(t) = -(4\pi t_L/L_0)\ln(1 - t/t_L)$.

Equation (4.12) suggests that a convenient choice of gauge coordinates $(\zeta^0, \zeta^1) = (u, v)$ for the trajectory of the evaporating ACO loop is

$$u = \frac{1}{2}(\phi(t) - \bar{\sigma}) \quad \text{and} \quad v = \frac{1}{2}(\phi(t) + \bar{\sigma}). \quad (4.13)$$

Then $1 - t/t_L = e^{-\kappa\mu(u+v)}$, where $\kappa\mu \equiv (4\pi t_L/L_0)^{-1}$ or equivalently $\kappa = \gamma^0/(4\pi)$. This in turn means that the time coordinate t on the loop can be written as $t = t_L[1 - e^{-\kappa\mu(u+v)}]$, and the world sheet $\mathbf{T}$ of the evaporating loop can be represented by the 4-function

$$X^a(u, v) = [t_L, \mathbf{0}]^a + \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)}[-\frac{1}{\kappa\mu}, \mathbf{r}(u, v)]^a \quad (4.14)$$

where

$$\mathbf{r}(u, v) = \cos(v)\hat{\mathbf{x}} + \sin(v)\hat{\mathbf{y}} + (|u| - \frac{1}{2}\pi)\hat{\mathbf{z}} \quad (4.15)$$

for $-\pi \leq u \leq \pi$, and $t_L = \frac{1}{4\pi\kappa\mu}L_0$.

Strictly speaking, the vector function $\mathbf{r}$ appearing in the 4-function (4.14) is the 2π -periodic extension (in u) of the function defined in (4.15). The gauge coordinate pairs (u, v) and $(u + 2\pi k, v - 2\pi k)$ therefore correspond to the same spacetime point for any integer k . However, for the sake of simplicity the range of the parameter u will henceforth be restricted to $(-\pi, \pi]$. The final evaporation point $x^a = [t_L, \mathbf{0}]^a$ of the loop then corresponds to the limit $v \rightarrow \infty$. Note that in this limit the scale factor $L(t) = L_0 e^{-\kappa\mu(u+v)}$ goes to zero, while the angular speed dv/dt of the loop diverges and $\mathbf{r}(u, v)$ is undefined (although the loop's 4-velocity X_t^a , being scale-free, remains subluminal everywhere).

If $|\kappa\mu(u + v)| \ll 1$ the loop's position function (4.14) can be expanded in powers of μ to give

$$X_{(0)}^a(u, v) = \frac{1}{4\pi} L_0 [u + v, \mathbf{r}(u, v)]^a \quad (4.16)$$

and

$$\delta X^a(u, v) = -\frac{1}{4\pi} L_0 \kappa\mu \left[\frac{1}{2}(u + v)^2, (u + v) \mathbf{r}(u, v) \right]^a. \quad (4.17)$$

It is these functions that need to satisfy the zeroth- and first-order equations of motion (2.13) and (2.14). Note that the leading-order position function (4.16) is effectively generated by an expansion of X^a about the curve $u + v = 0$ on the loop's world sheet, or equivalently about the $t = 0$ spacelike cross-section.

The position function X^a can be expanded about other cross-sections of constant $t < t_L$ by writing $v = v_* + \bar{v}$ where v_* is any fixed real number. Then equation (4.16) becomes

$$X^a(u, \bar{v}) = [t_L, \mathbf{0}]^a + \frac{1}{4\pi} L_* e^{-\kappa\mu(u+\bar{v})} \left[-\frac{1}{\kappa\mu}, \mathbf{r}(u, v_* + \bar{v}) \right]^a \quad (4.18)$$

with $L_* \equiv L_0 e^{-\kappa\mu v_*}$, and if $|\kappa\mu(u + \bar{v})| \ll 1$ the zeroth-order position function is

$$X_{(0)}^a(u, \bar{v}) = [t_*, \mathbf{0}]^a + \frac{1}{4\pi} L_* [u + \bar{v}, \mathbf{r}(u, v_* + \bar{v})]^a \quad (4.19)$$

where $t_* \equiv \frac{1}{4\pi\kappa\mu} L_0 (1 - e^{-\kappa\mu v_*})$. The analogue of equation (4.17) for δX^a is found by replacing L_0 with L_* , $u + v$ with $u + \bar{v}$, and $\mathbf{r}(u, v)$ with $\mathbf{r}(u, v_* + \bar{v})$. Clearly, any two expansions of this type will be related by a time translation, a constant dilation, and a rotation in the x - y plane, and if any of the expansions satisfies the equations of motion (2.13) and (2.14) then they all will.

The equations of motion will be considered more closely in Section 6. It will come as no surprise that the zeroth-order position function (4.16) does satisfy the zeroth-order equation of motion (2.13), which reads simply $X_{(0),uv}^a = 0$. Before considering the first-order equation of motion (2.14), however, it is necessary to

calculate the metric perturbations h_{ab} , which depend only on $X_{(0)}^a$. This will be done in Section 5.

After a long calculation described in detail in Section 6, it turns out that the first-order position function δX^a given by (4.17) does *not* satisfy the first-order equation of motion (2.14). The reason for this is not difficult to appreciate. As was mentioned at the beginning of this section, the calculation leading to the evaporating loop trajectory (4.14) was predicated on the assumption that the background metric is flat. It is evident from (2.14), however, that the first-order equation of motion involves the first-order metric perturbation h_{ab} . Once the background metric g_{ab} ceases to be flat, the spacetime coordinates $[t, x, y, z]$ no longer measure proper times or proper lengths. In particular, the self-similar shrinkage driven by the scale factor $L_0 e^{-\kappa\mu(u+v)}$ may proceed at different rates in different directions at various points on the world sheet $\mathbf{T}$.

This possibility is easily accounted for by adding scale corrections, of order μ , to the components of the evaporating loop's position function (4.14). The form of these scale corrections is of course constrained by the equation of motion (2.14), but even so there remains considerable freedom in the choice of the correction functions, corresponding not only to possible transformations (at order μ) of the gauge coordinates (u, v) and the spacetime coordinates $[t, x, y, z]$, but also to the propagation of free vibrations (with amplitude μ) around the loop.

In what follows, the freedom in the choice of gauge coordinates will be partially fixed by taking the position vector to have the form

$$X^a(u, v) = [t_L, \mathbf{0}]^a + \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} \left[-\frac{1}{\kappa\mu} e^{\kappa\mu^2 F(u,v)}, \bar{\mathbf{r}}(u, v) \right]^a, \quad (4.20)$$

where now

$$\bar{\mathbf{r}}(u, v) = e^{\mu G(u,v)} [\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}] + [(|u| - \frac{1}{2}\pi) + \mu A(u, v)] \hat{\mathbf{z}} \quad (4.21)$$

for $-\pi \leq u \leq \pi$. Here, the correction functions F , G and A are assumed to be continuous and 2π -periodic in both u and v , so that (u, v) and $(u + 2\pi, v - 2\pi)$ again correspond to the same spacetime point, and all three functions remain small compared to μ^{-1} as $u + v \rightarrow \infty$. The assumption that the trajectory is dilated by the same factor $e^{\mu G}$ in the x and y directions effectively removes any freedom to redefine the gauge coordinate v .³

³To see this, suppose that the term $G(u, v)\{\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}\}$ in equation (4.23) has the more general form $G(u, v) \cos(v) \hat{\mathbf{x}} + H(u, v) \sin(v) \hat{\mathbf{y}}$. Then if v is replaced by a new coordinate v' defined by $v = v' + \mu V(u, v')$ for some function V , the correction term $G \cos(v) \hat{\mathbf{x}} + H \sin(v) \hat{\mathbf{y}}$

The presence of free vibrations which break the self-similarity of the world-sheet can be removed by requiring the intrinsic 2-metric $\gamma_{AB} = g_{ab}X^a_{,A}X^b_{,B}$ to have the strictly self-similar form

$$\gamma_{AB}(u, v) = (\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)} \bar{\gamma}_{AB}(u) \quad (4.22)$$

to linear order in μ , with $\bar{\gamma}_{AB}$ a 2-metric to be determined. This constraint will be considered in more detail in Section 6.2.

Given (4.20), the leading-order position function $X^a_{(0)}$ near $t = 0$ retains the form (4.16), while the first-order position function becomes

$$\begin{aligned} \delta X^a(u, v) = & -\frac{1}{4\pi}L_0\kappa\mu[\frac{1}{2}(u+v)^2, (u+v)\mathbf{r}(u, v)]^a \\ & +\frac{1}{4\pi}L_0\mu[-F(u, v), G(u, v)\{\cos(v)\hat{\mathbf{x}} + \sin(v)\hat{\mathbf{y}}\} + A(u, v)\hat{\mathbf{z}}]^a. \end{aligned} \quad (4.23)$$

The actual form of the correction functions F , G and A will be determined once the first-order equation of motion (2.14) has been solved, and additional gauge conditions have been imposed.

Two remarks about the final form (4.20) of the loop trajectory are in order here. The first is that although (4.20) was developed to describe an evaporating ACO loop over the time interval $0 \leq t < t_L$ (or equivalently $0 \leq u + v < \infty$), it can without difficulty be assumed to extend over the entire interval $-\infty < t < t_L$ (or $-\infty < u + v < \infty$). The trajectory then describes a loop that has been evaporating “eternally”, with its energy and length scaling (to leading order in μ) as $1 - t/t_L$. This is obviously just a mathematical idealization, but it does side-step the need to construct initial data for the metric perturbations h_{ab} at $t = 0$.

The second point is that the value of the constant κ appearing in the corrected trajectory (4.20) is initially undetermined. Although it is to be expected in view of the results of Paper I that $\kappa \approx 39.0025/(4\pi) \approx 3.1037$, no particular value of κ is assumed in the analysis that follows. It is only after the first-order equation of motion is solved that it becomes evident that the periodicity constraint on G forces κ to take on its expected value.

A schematic representation of the evaporation of the loop is shown in Figure 2. The thickened line corresponds to the outer envelope of the loop, whose radius

becomes $(G \cos v' - V \sin v')\hat{\mathbf{x}} + (H \sin v' + V \cos v')\hat{\mathbf{y}}$ to leading order in μ . The unique gauge choice $V = (G - H) \sin v' \cos v'$ then reduces this term to the isotropic form $G'\{\cos(v')\hat{\mathbf{x}} + \sin(v')\hat{\mathbf{y}}\}$, with $G' = G \cos^2 v' + H \sin^2 v'$.

shrinks to zero at the final evaporation point $x^a = [t_L, \mathbf{0}]^a$. Also shown are the future light cones F_0 and F_L of the origin and the evaporation point respectively. Note that, because of the first-order corrections to the background metric and the loop trajectory X^a , the envelope of the loop and the light cones depicted in Figure 2 should deviate from straight lines by terms of order μ .

Now that the choice of the evaporating loop trajectory (4.20) has been explained in some depth, the remainder of the paper is a straightforward application of the solution methods discussed in Section 2. The metric perturbations h_{ab} generated by $X_{(0)}^a$ are calculated in Section 5, the equation of motion for the perturbation functions δX^a is constructed and solved in Section 6, and the metric perturbations on the future light cone F_L of the final evaporation point are shown to match onto an empty, flat spacetime in Section 7.

5. The Metric Perturbations

5.1. Rotating self-similarity of the metric perturbations

The metric perturbations h_{ab} at any field point $x^a \equiv [t, \mathbf{x}]^a$ away from the world sheet of the evaporating loop can be calculated from equation (2.10). Because the analysis of Section 7 depends on an understanding of the behavior of h_{ab} arbitrarily close to the light cone F_L , it will be necessary to consider source points on the evaporating loop close to the final evaporation point (where $u + v \rightarrow \infty$) and so it cannot be assumed that $|\kappa\mu(u + v)| \ll 1$. The scale factor $L_0 e^{-\kappa\mu(u+v)}$ will therefore be retained in full in this section, although the correction functions F , G and A – which are bounded and appear only at order μ in (4.20) – will be ignored. In other words, the trajectory will in this section be assumed to have its flat-space form (4.14).

The first step in the calculation is to write down the leading-order expressions for the intrinsic 2-metric γ_{AB} and the source function S_{ab} . Since

$$X^a{}_{,u} = \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} [1, \text{sgn}(u) \hat{\mathbf{z}}]^a \quad (5.1)$$

and

$$X^a{}_{,v} = \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} [1, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}}]^a \quad (5.2)$$

to leading order in μ , it follows that the intrinsic 2-metric $\gamma_{AB} = X_{,A} \cdot X_{,B}$ on the world sheet has components

$$\gamma_{uu} = \gamma_{vv} = 0 \quad \text{and} \quad \gamma_{uv} = \left(\frac{1}{4\pi} L_0\right)^2 e^{-2\kappa\mu(u+v)} \quad (5.3)$$

to the same order. Hence, $\gamma^{1/2} = (\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)}$, while the inverse 2-metric γ^{AB} has components

$$\gamma^{uu} = \gamma^{vv} = 0 \quad \text{and} \quad \gamma^{uv} = (\frac{1}{4\pi}L_0)^{-2} e^{2\kappa\mu(u+v)}. \quad (5.4)$$

The source function defined in equation (2.11) can be written as

$$S_{ab}(x'^d) = \mu \int \Psi_{ab}(u, v) \delta^{(4)}(x'^d - X^d) du dv \quad (5.5)$$

where

$$\Psi_{ab} \equiv \gamma^{1/2} \gamma^{AB} (X_{a,A} X_{b,B} - \frac{1}{2} \eta_{ab} X_{,A} \cdot X_{,B}), \quad (5.6)$$

evaluated again at leading order in μ . Note here that $\gamma^{AB} X_{a,A} X_{b,B}$ is just the world-sheet projection tensor p_{ab} , which in matrix form is

$$p_{ab} = \begin{bmatrix} 2 & \sin v & -\cos v & -s \\ \sin v & 0 & 0 & -s \sin v \\ -\cos v & 0 & 0 & s \cos v \\ -s & -s \sin v & s \cos v & 0 \end{bmatrix}_{ab} \quad (5.7)$$

where $s \equiv \text{sgn}(u)$. The function Ψ_{ab} is therefore equal to $-\gamma^{1/2} q_{ab}$, where $q_{ab} \equiv \eta_{ab} - p_{ab}$ was introduced in Section 2 as the orthogonal complement of p_{ab} . Hence, $\Psi_{ab} = (\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)} \bar{\Psi}_{ab}$ with

$$\bar{\Psi}_{ab}(s, v) \equiv \begin{bmatrix} 1 & \sin v & -\cos v & -s \\ \sin v & 1 & 0 & -s \sin v \\ -\cos v & 0 & 1 & s \cos v \\ -s & -s \sin v & s \cos v & 1 \end{bmatrix}_{ab}. \quad (5.8)$$

Given (5.5), the integral expression (2.10) for h_{ab} reads

$$h_{ab}(t, \mathbf{x}) = -4\mu \int \frac{d^3 x'}{|\mathbf{x} - \mathbf{x}'|} \int \Psi_{ab}(u, v) \delta^{(4)}(x'^d - X^d) du dv \quad (5.9)$$

where the source point is $x'^d = [t', \mathbf{x}']^d$, with $t' \equiv t - |\mathbf{x} - \mathbf{x}'|$ the retarded time⁴. The distributional factor $\delta^{(4)}(x'^d - X^d)$ in (5.9) can be integrated out by first

⁴Note that, at a physical level, the equation $t' = t - |\mathbf{x} - \mathbf{x}'|$ imposes the requirement that the source point $[t', \mathbf{x}']$ lie on the backwards light cone of the field point $[t, \mathbf{x}]$ to leading order in μ .

transforming from $y^a \equiv [u, \mathbf{x}']^a$ to $z^a \equiv [t', \mathbf{x}']^a - X^a(u, v)$, with v , t and $\mathbf{x}$ held constant. The Jacobian of this transformation is:

$$|\partial y^a / \partial z^b| = |\partial z^b / \partial y^a|^{-1} = |X^0_{,u} - \mathbf{n} \cdot \mathbf{X}_{,u}|^{-1}, \quad (5.10)$$

where $\mathbf{n} = (\mathbf{x} - \mathbf{x}')/|\mathbf{x} - \mathbf{x}'|$ is the unit vector from the source point $\mathbf{x}'$ to the field point $\mathbf{x}$.

On integrating over z^a , t' is everywhere replaced by $X^0 \equiv t_L[1 - e^{-\kappa\mu(u+v)}]$ and $\mathbf{x}'$ by $\mathbf{X} \equiv \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)}\mathbf{r}(u, v)$, while u becomes an implicit function of v , t and $\mathbf{x}$ through the equation $X^0(u, v) = t - |\mathbf{x} - \mathbf{X}(u, v)|$. The integral for h_{ab} therefore reads

$$h_{ab}(t, \mathbf{x}) = -4\mu \int ||\mathbf{x} - \mathbf{X}| X^0_{,u} - (\mathbf{x} - \mathbf{X}) \cdot \mathbf{X}_{,u}|^{-1} \Psi_{ab}(u, v) dv. \quad (5.11)$$

where, as before, $X^0_{,u} = \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)}$ and $\mathbf{X}_{,u} = \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)}s \hat{\mathbf{z}}$ to leading order in μ .

Evaluation of the integral (5.11) is simplified by the fact that h_{ab} satisfies a very elegant self-similarity relation. If $[t, \mathbf{x}]$ is any field point with $t - t_L < |\mathbf{x}|$ (where $t - t_L = |\mathbf{x}|$ is just the equation, at leading order, of the future light cone F_L of the final evaporation point), first define

$$\psi(t, \mathbf{x}) \equiv -(\kappa\mu)^{-1} \ln[(|\mathbf{x}| - t + t_L)/t_L] \quad (5.12)$$

then consider a second field point $[\bar{t}, \bar{\mathbf{x}}]$ with

$$\bar{t} \equiv e^{\kappa\mu\psi} |\mathbf{x}| \quad \text{and} \quad \bar{\mathbf{x}} \equiv e^{\kappa\mu\psi} R(-\psi) \mathbf{x} \quad (5.13)$$

where

$$R(\theta) \equiv \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (5.14)$$

is the standard rotation matrix about the z -axis. Note that $[\bar{t}, \bar{\mathbf{x}}]$ lies on the future light cone F_0 of the origin, as $\bar{t} = |\bar{\mathbf{x}}|$.

Now, the equation $t - X^0(u, v) = |\mathbf{x} - \mathbf{X}(u, v)|$ which specifies u as a function of v in the integral (5.11) – or equivalently on the backwards light cone of the original field point $[t, \mathbf{x}]$ – reads explicitly

$$t - t_L[1 - e^{-\kappa\mu(u+v)}] = |\mathbf{x} - \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)}\mathbf{r}(u, v)|. \quad (5.15)$$

Multiplying this equation by $e^{\kappa\mu\psi(t,\mathbf{x})}$ gives:

$$(t - t_L)e^{\kappa\mu\psi} + t_L e^{-\kappa\mu(u+v-\psi)} = |e^{\kappa\mu\psi} \mathbf{x} - \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v-\psi)} \mathbf{r}(u, v)| \quad (5.16)$$

where $(t - t_L)e^{\kappa\mu\psi} = e^{\kappa\mu\psi} |\mathbf{x}| - t_L$ and

$$\begin{aligned} \mathbf{r}(u, v) &\equiv \cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}} + (|u| - \frac{1}{2}\pi) \hat{\mathbf{z}} \\ &= R(\psi) [\cos(v - \psi) \hat{\mathbf{x}} + \sin(v - \psi) \hat{\mathbf{y}} + (|u| - \frac{1}{2}\pi) \hat{\mathbf{z}}] \\ &\equiv R(\psi) \mathbf{r}(u, v - \psi). \end{aligned} \quad (5.17)$$

Since $|\mathbf{v}| = |R(-\psi)\mathbf{v}|$ for any 3-vector $\mathbf{v}$, and $R(-\psi)R(\psi) = I$, equation (5.16) can be rewritten as

$$\bar{t} - t_L [1 - e^{-\kappa\mu(u+v-\psi)}] = |\bar{\mathbf{x}} - \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v-\psi)} \mathbf{r}(u, v - \psi)|, \quad (5.18)$$

or equivalently $\bar{t} - X^0(u, v - \psi) = |\bar{\mathbf{x}} - \mathbf{X}(u, v - \psi)|$.

Hence, if $u = U(v)$ is the equation of the curve of intersection of the world sheet $\mathbf{T}$ with the backwards light cone of a given field point $[t, \mathbf{x}]$, then the equation for u on the backwards light cone of the corresponding image point $[\bar{t}, \bar{\mathbf{x}}]$ on F_0 is $u = U(v - \psi(t, \mathbf{x}))$. Similarly, the integrand in equation (5.11) for h_{ab} , which is equivalent to

$$\frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} ||\mathbf{x} - \mathbf{X}(u, v)| - s [\mathbf{x} - \mathbf{X}(u, v)] \cdot \hat{\mathbf{z}}|^{-1} \bar{\Psi}_{ab}(s, v), \quad (5.19)$$

can be re-expressed as⁵

$$\begin{aligned} &\frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v-\psi)} ||e^{\kappa\mu\psi} R(-\psi) \mathbf{x} - \mathbf{X}(u, v - \psi)| - s [e^{\kappa\mu\psi} R(-\psi) \mathbf{x} - \mathbf{X}(u, v - \psi)] \cdot \hat{\mathbf{z}}|^{-1} \\ &\times \bar{\Psi}_{ab}(s, v) \\ &= \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v-\psi)} ||\bar{\mathbf{x}} - \mathbf{X}(u, v - \psi)| - s [\bar{\mathbf{x}} - \mathbf{X}(u, v - \psi)] \cdot \hat{\mathbf{z}}|^{-1} \\ &\times R_a^c(\psi) \bar{\Psi}_{cd}(s, v - \psi) R_b^d(\psi) \end{aligned} \quad (5.20)$$

where

$$R_a^c(\psi) \equiv (1 \oplus R(\psi))_a^c = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \psi & -\sin \psi & 0 \\ 0 & \sin \psi & \cos \psi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \quad (5.21)$$

⁵Note here that

$$e^{\kappa\mu\psi} [\mathbf{x} - \mathbf{X}(u, v)] \cdot \hat{\mathbf{z}} = [e^{\kappa\mu\psi} R(-\psi) \mathbf{x} - \mathbf{X}(u, v - \psi)] \cdot \hat{\mathbf{z}}$$

for the simple reason that the z -component of $\mathbf{x}$ is unaffected by the rotation matrix R , while the z -component of $\mathbf{X}$ is $\frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} (|u| - \frac{1}{2}\pi)$.

Comparison of equations (5.19) and (5.20), together with the fact that if $u = U(v)$ on the backwards light cone of $[t, \mathbf{x}]$ then $u = U(v - \psi)$ on the backwards light cone of $[\bar{t}, \bar{\mathbf{x}}]$, indicates that

$$h_{ab}(t, \mathbf{x}) = R_a^c(\psi) h_{cd}(\bar{t}, \bar{\mathbf{x}}) R_b^d(\psi), \quad (5.22)$$

where $\psi(t, \mathbf{x})$ is given by equation (5.12). The geometric meaning of this relation is illustrated in Figure 3. To calculate h_{ab} at a field point $[t, \mathbf{x}]$ in the past of F_L , simply dilate the point by a factor $e^{\kappa\mu\psi}$ through $[t_L, \mathbf{0}]$ and rotate it by an angle $-\psi$ in the x - y plane to map it to the image point $[\bar{t}, \bar{\mathbf{x}}]$ on F_0 . Then calculate h_{ab} at $[\bar{t}, \bar{\mathbf{x}}]$ and rotate each of the components of the resulting tensor by an angle ψ in the x - y plane to give h_{ab} at $[t, \mathbf{x}]$.

From a computational viewpoint, equation (5.22) allows us to calculate h_{ab} at any point away from the world sheet of the loop from a knowledge of h_{ab} on the null surface F_0 alone. Furthermore, it points to the existence of a symmetry in the background metric induced by the self-similarity of the loop's trajectory. This symmetry can be better appreciated by writing the components (5.13) of the image point explicitly in terms of t and $\mathbf{x}$:

$$\bar{t} \equiv t_L(1 + |\mathbf{w}|)^{-1}|\mathbf{w}| \quad \text{and} \quad \bar{\mathbf{x}} \equiv t_L(1 + |\mathbf{w}|)^{-1}R(-\psi)\mathbf{w} \quad (5.23)$$

where

$$\mathbf{w} \equiv \mathbf{x}/(t_L - t). \quad (5.24)$$

If it were not for the presence of the rotation operators in (5.22) and (5.23), which depend on ψ rather than $\mathbf{w}$, the metric $\eta_{ab} + h_{ab}$ would be a function of $\mathbf{w}$ alone and so would be strictly self-similar. As it is, the rotation operators act to enforce a slightly more complicated symmetry in the metric which could perhaps be termed “rotating self-similarity”.

5.2. Explicit form of the metric perturbations on F_0

Suppose now that $[[\bar{\mathbf{x}}], \bar{\mathbf{x}}]$ is a general point on the future light cone F_0 of the origin. The equation (5.15) for u as a function of v on the backwards light cone of $[[\bar{\mathbf{x}}], \bar{\mathbf{x}}]$ reads:

$$|\bar{\mathbf{x}}| - t_L[1 - e^{-\kappa\mu(u+v)}] = |\bar{\mathbf{x}} - \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)} \mathbf{r}(u, v)|. \quad (5.25)$$

Application of the triangle inequality to the norm on the right-hand side of this equation then gives:

$$t_L|1 - e^{-\kappa\mu(u+v)}| \leq \frac{1}{4\pi}L_0 e^{-\kappa\mu(u+v)}[1 + (|u| - \frac{1}{2}\pi)^2]^{1/2}, \quad (5.26)$$

as $|\mathbf{r}(u, v)| = [1 + (|u| - \frac{1}{2}\pi)^2]^{1/2}$. Since $t_L = \frac{1}{4\pi\kappa\mu}L_0$ and the maximum value of $[1 + (|u| - \frac{1}{2}\pi)^2]^{1/2}$ for $u \in (-\pi, \pi]$ is $(1 + \frac{1}{4}\pi^2)^{1/2} \approx 1.8621$, this inequality can be re-expressed as

$$|e^{\kappa\mu(u+v)} - 1| \leq (1 + \frac{1}{4}\pi^2)^{1/2} \kappa\mu. \quad (5.27)$$

It is therefore evident that $|\kappa\mu(u+v)|$ will remain small at all points on the backwards light cone of $[[\bar{\mathbf{x}}, \bar{\mathbf{x}}]$ provided that $\kappa\mu$ itself is sufficiently small. For example, $|\kappa\mu(u+v)|$ will be everywhere no greater than 10^{-2} if $\ln[1 - (1 + \frac{1}{4}\pi^2)^{1/2} \kappa\mu] \geq -10^{-2}$, and this in turn will be true if $\kappa\mu \lesssim 5.34 \times 10^{-3}$. Given that we expect that $\kappa \approx 3.1037$ and μ will be of order 10^{-6} or smaller for a GUT string, the assumption that $|\kappa\mu(u+v)|$ remains small is not an unreasonable one.

Expanding $e^{-\kappa\mu(u+v)}$ in powers of μ in (5.25) and retaining only the leading-order terms gives the following equation for u as a function of v :

$$T - (u + v) = [(X - \cos v)^2 + (Y - \sin v)^2 + \{Z - (|u| - \frac{1}{2}\pi)\}^2]^{1/2}, \quad (5.28)$$

where

$$(X, Y, Z) \equiv 4\pi L_0^{-1}(\bar{x}, \bar{y}, \bar{z}) \quad \text{and} \quad T \equiv (X^2 + Y^2 + Z^2)^{1/2}. \quad (5.29)$$

Equation (5.28) can easily be solved to give an explicit formula for u , but it turns out that in calculating h_{ab} it is more useful to regard u as the independent variable.

Similarly, for small values of $|\kappa\mu(u+v)|$ the integrand (5.19) in the formula (5.11) for h_{ab} at $[[\bar{\mathbf{x}}, \bar{\mathbf{x}}]$ becomes:

$$\{[(X - \cos v)^2 + (Y - \sin v)^2 + \{Z - (|u| - \frac{1}{2}\pi)\}^2]^{1/2} - s[Z - (|u| - \frac{1}{2}\pi)]\}^{-1} \bar{\Psi}_{ab}(s, v). \quad (5.30)$$

Note that the denominator in this expression is manifestly positive or zero. In view of equation (5.28) and the fact that $s|u| = u$, the denominator can be expressed in the simpler form

$$[(X - \cos v)^2 + (Y - \sin v)^2 + \{Z - (|u| - \frac{1}{2}\pi)\}^2]^{1/2} - s[Z - (|u| - \frac{1}{2}\pi)] = T - s(Z + \frac{1}{2}\pi) - v, \quad (5.31)$$

and (5.11) reduces to

$$h_{ab}([\bar{\mathbf{x}}, \bar{\mathbf{x}}]) = -4\mu \int [T - s(Z + \frac{1}{2}\pi) - v]^{-1} \bar{\Psi}_{ab}(s, v) dv. \quad (5.32)$$

All that remains now is to determine the limits of integration. The entire curve of intersection Γ of the world sheet with the backwards light cone of $[[\bar{\mathbf{x}}, \bar{\mathbf{x}}]$ will

be traced out as u varies from $-\pi$ to π . The integral (5.32) therefore breaks into two parts: one with $s = -1$ and $u \in [-\pi, 0]$, and one with $s = +1$ and $u \in [0, \pi]$. The corresponding values of v , which I will denote as $V_k \equiv v|_{u=k\pi}$ for $k = -1, 0$ and 1 , are roots of the light-cone equation (5.28):

$$T - (k\pi + V_k) = [(X - \cos V_k)^2 + (Y - \sin V_k)^2 + \{Z - (|k| - \frac{1}{2})\pi\}^2]^{1/2}. \quad (5.33)$$

Although it is not possible to solve for V_k explicitly except in certain special cases, it is clear by inspection that $V_{-1} = V_1 + 2\pi$. Furthermore, implicit differentiation of (5.28) with $[T, X, Y, Z]$ constant indicates that

$$\frac{dv}{du} = -\frac{T - s(Z + \frac{1}{2}\pi) - v}{T - (u + v) + X \sin v - Y \cos v}. \quad (5.34)$$

Some of the mathematical consequences of equations (5.33) and (5.34) are examined in Appendix A. In particular, it is shown that the values of V_0 and V_1 are bounded on F_0 , and that v is a monotonic decreasing function of u . The last property follows partly from the fact that the numerator in (5.34) is positive or zero from (5.31), while the denominator is positive or zero as a consequence of (5.28) and the identity

$$(X - \cos v)^2 + (Y - \sin v)^2 = (X \sin v - Y \cos v)^2 + (X \cos v + Y \sin v - 1)^2. \quad (5.35)$$

However, cases where either the numerator or denominator is zero for some value of u need to be considered separately, and it is shown in Appendices A.2 and A.3 that the corresponding spacetime points have interesting physical interpretations. Perhaps the most important result is that the spacetime derivatives of h_{ab} are undefined at all points with the property that the denominator in (5.34) is zero at one of the extreme values $u = 0$ or $\pm\pi$.

Given that $V_{-1} = V_1 + 2\pi \geq V_0 \geq V_1$, the equation for the metric perturbations becomes:

$$\begin{aligned} h_{ab}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}}) &= -4\mu \int_{V_0}^{V_1+2\pi} [T + (Z + \frac{1}{2}\pi) - v]^{-1} \bar{\Psi}_{ab}(-1, v) dv \\ &\quad - 4\mu \int_{V_1}^{V_0} [T - (Z + \frac{1}{2}\pi) - v]^{-1} \bar{\Psi}_{ab}(+1, v) dv. \end{aligned} \quad (5.36)$$

After referring back to the definition (5.8) of $\bar{\Psi}_{ab}$, it is readily seen that

$$h_{ab}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}}) = 4\mu \begin{bmatrix} I_+ + I_- & S_+ + S_- & -C_+ - C_- & I_+ - I_- \\ S_+ + S_- & I_+ + I_- & 0 & S_+ - S_- \\ -C_+ - C_- & 0 & I_+ + I_- & -C_+ + C_- \\ I_+ - I_- & S_+ - S_- & -C_+ + C_- & I_+ + I_- \end{bmatrix}_{ab} \quad (5.37)$$

where

$$I_+ = \ln[(\chi_+ - V_1 - 2\pi)/(\chi_+ - V_0)], \quad I_- = \ln[(\chi_- - V_0)/(\chi_- - V_1)], \quad (5.38)$$

$$\begin{bmatrix} S_+ \\ C_+ \end{bmatrix} = \begin{bmatrix} -\cos \chi_+ & \sin \chi_+ \\ \sin \chi_+ & \cos \chi_+ \end{bmatrix} \begin{bmatrix} \text{Si}(\chi_+ - V_1 - 2\pi) - \text{Si}(\chi_+ - V_0) \\ \text{Ci}(\chi_+ - V_1 - 2\pi) - \text{Ci}(\chi_+ - V_0) \end{bmatrix} \quad (5.39)$$

and

$$\begin{bmatrix} S_- \\ C_- \end{bmatrix} = \begin{bmatrix} -\cos \chi_- & \sin \chi_- \\ \sin \chi_- & \cos \chi_- \end{bmatrix} \begin{bmatrix} \text{Si}(\chi_- - V_0) - \text{Si}(\chi_- - V_1) \\ \text{Ci}(\chi_- - V_0) - \text{Ci}(\chi_- - V_1) \end{bmatrix}. \quad (5.40)$$

with $\chi_{\pm} \equiv T \pm (Z + \frac{\pi}{2})$, and $\text{Si}(x) = \int_0^x w^{-1} \sin w \, dw$ and $\text{Ci}(x) = -\int_x^\infty w^{-1} \cos w \, dw$ the usual sine and cosine integrals.

One case in which it is possible to solve explicitly for V_0 and V_1 , and so to write explicit formulas for the metric perturbations h_{ab} , occurs when the image point $[|\bar{\mathbf{x}}|, \bar{\mathbf{x}}]$ lies on the $\bar{z}$ -axis. Then $X = Y = 0$ and $T = |Z|$, and equation (5.33) reduces to

$$V_k = |Z| - k\pi - [1 + \{Z - (|k| - \frac{1}{2})\pi\}^2]^{1/2}. \quad (5.41)$$

In particular,

$$V_0 = |Z| - [1 + (Z + \frac{\pi}{2})^2]^{1/2} \quad \text{and} \quad V_1 = |Z| - \pi - [1 + (Z - \frac{\pi}{2})^2]^{1/2} \quad (5.42)$$

while $\chi_{\pm} = |Z| \pm (Z + \frac{\pi}{2})$.

Hence,

$$\begin{aligned} & (\chi_+ - V_1 - 2\pi)/(\chi_+ - V_0) \\ &= (\chi_- - V_0)/(\chi_- - V_1) \\ &= \{Z - \frac{\pi}{2} + [1 + (Z - \frac{\pi}{2})^2]^{1/2}\} \{-Z - \frac{\pi}{2} + [1 + (Z + \frac{\pi}{2})^2]^{1/2}\} \end{aligned} \quad (5.43)$$

and $I_- = I_+$. This means that $h_{tz} = 0$, and that

$$\begin{aligned} & h_{tt} = h_{xx} = h_{yy} = h_{zz} \\ &= 8\mu \ln(\{Z - \frac{\pi}{2} + [1 + (Z - \frac{\pi}{2})^2]^{1/2}\} \{-Z - \frac{\pi}{2} + [1 + (Z + \frac{\pi}{2})^2]^{1/2}\}) \end{aligned} \quad (5.44)$$

increases monotonically from approximately -19.734μ (at $Z = 0$) to 0 (as $|Z| \rightarrow \infty$). In fact, for large values of $|Z|$,

$$h_{tt} = h_{xx} = h_{yy} = h_{zz} = -8\pi\mu/|Z| + O(|Z|^{-2}). \quad (5.45)$$

The variation of $h_{tt}/(4\mu)$ with Z is plotted in Figure 4a.

Explicit formulas for the remaining non-zero metric perturbations, $h_{tx} = 4\mu(S_+ + S_-)$, $h_{ty} = -4\mu(C_+ + C_-)$, $h_{zx} = 4\mu(S_+ - S_-)$ and $h_{zy} = -4\mu(C_+ - C_-)$ can be found by substituting the expressions (5.42) into (5.39) and (5.40). However, there is little to be gained from writing these formulas out in full. It is easily seen that if Z is replaced by $-Z$ then $S_+ \leftrightarrow -S_-$ and $C_+ \leftrightarrow -C_-$, and so $h_{tx} \rightarrow -h_{tx}$, $h_{ty} \rightarrow -h_{ty}$, $h_{zx} \rightarrow h_{zx}$ and $h_{zy} \rightarrow h_{zy}$. This set of symmetries is to be expected on physical grounds, as the ACO loop is invariant under a complete spatial inversion ($x \rightarrow -x$, $y \rightarrow -y$ and $z \rightarrow -z$), and therefore in this special case the space-space components of h_{ab} should be even functions of Z , and the time-space components odd functions of Z .

The variations of $h_{tx}/(4\mu)$, $h_{ty}/(4\mu)$, $h_{zx}/(4\mu)$ and $h_{zy}/(4\mu)$ are plotted in Figures 4b to 4e respectively. At $Z = 0$, the perturbations h_{tx} and h_{ty} are of course both zero, while $h_{zx} \approx -5.308\mu$ and $h_{zy} \approx 12.395\mu$. All the perturbations tend to zero as $|Z| \rightarrow \infty$, with

$$h_{tx} = 4\pi\mu/Z + O(|Z|^{-3}), \quad (5.46)$$

$$h_{ty} = -4\pi\mu/(Z|Z|) + O(|Z|^{-4}), \quad (5.47)$$

$$h_{zx} = -4\pi\mu/|Z| + O(|Z|^{-3}) \quad (5.48)$$

and

$$h_{zy} = \frac{13}{6}\pi\mu/Z^4 + O(|Z|^{-6}). \quad (5.49)$$

6. Solving the String Equations of Motion

6.1. Near-field form of the metric perturbations

For the purposes of setting out the first-order equation of motion (2.14), it is necessary to generate expressions for the metric perturbations h_{ab} in the neighborhood of the world sheet of the evaporating loop. As was mentioned in Section 4, when doing this it is sufficient to restrict attention to points near the spacelike surface $t = 0$ only. So consider a field point $x^a = [t, \mathbf{x}]^a$ of the form $X^a(u, v) + \delta x^a$, where $X^a(u, v)$ is a fixed point on the flat-space trajectory (4.14) with $|\kappa\mu(u + v)| \ll 1$, and the components of the displacement δx^a are all of the order of some small parameter.

A first point to note is that if $X^a(\bar{u}, \bar{v})$ is any source point on the loop's trajectory on the backwards light cone of $X^a(u, v)$ then $|\kappa\mu(\bar{u} + \bar{v})|$ will also be

small compared to 1 for reasonable values of $\kappa\mu$. This follows from equation (5.15), which in the current situation reads:

$$t_L[1 - e^{-\kappa\mu(u+v)}] - t_L[1 - e^{-\kappa\mu(\bar{u}+\bar{v})}] = \left| \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} \mathbf{r}(u, v) - \frac{1}{4\pi} L_0 e^{-\kappa\mu(\bar{u}+\bar{v})} \mathbf{r}(\bar{u}, \bar{v}) \right|. \quad (6.1)$$

An application of the triangle inequality gives

$$t_L |e^{-\kappa\mu(\bar{u}+\bar{v})} - e^{-\kappa\mu(u+v)}| \leq \frac{1}{4\pi} L_0 e^{-\kappa\mu(u+v)} |\mathbf{r}(u, v)| + \frac{1}{4\pi} L_0 e^{-\kappa\mu(\bar{u}+\bar{v})} |\mathbf{r}(\bar{u}, \bar{v})|, \quad (6.2)$$

or equivalently (in parallel with the derivation of (5.27)),

$$|1 - e^{\kappa\mu(\bar{u}+\bar{v}) - \kappa\mu(u+v)}| \leq (1 + e^{\kappa\mu(\bar{u}+\bar{v}) - \kappa\mu(u+v)}) (1 + \frac{1}{4}\pi^2)^{1/2} \kappa\mu. \quad (6.3)$$

Hence,

$$|\kappa\mu(\bar{u} + \bar{v}) - \kappa\mu(u + v)| \leq \ln \left[\frac{1 + (1 + \frac{1}{4}\pi^2)^{1/2} \kappa\mu}{1 - (1 + \frac{1}{4}\pi^2)^{1/2} \kappa\mu} \right], \quad (6.4)$$

and so, for example, $|\kappa\mu(\bar{u} + \bar{v})|$ will be no greater than 10^{-2} everywhere if $\kappa\mu \lesssim 2.69 \times 10^{-3}$.

Since $|\kappa\mu(\bar{u} + \bar{v})|$ is guaranteed to be small, the equations (5.33) for V_k and (5.37)-(5.40) for h_{ab} continue to apply in this case, with

$$T = u + v + 4\pi L_0^{-1} \delta t \quad \text{and} \quad (X, Y, Z)^T = \mathbf{r}(u, v) + 4\pi L_0^{-1} \delta \mathbf{x}. \quad (6.5)$$

where as before $\mathbf{r}(u, v) = \cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}} + (|u| - \frac{1}{2}\pi) \hat{\mathbf{z}}$. In particular, if $\delta x^a = 0$ the equation for V_k becomes:

$$u + v - (k\pi + V_k) = [2 - 2\cos(v - V_k) + (|u| - |k|\pi)^2]^{1/2}, \quad (6.6)$$

and, given that u is restricted to the range $(-\pi, \pi]$, the trivial solution $V_k = v$ holds in two cases: (i) if $u > 0$ and $k = 0$; and (ii) if $u < 0$ and $k = -1$.

In the analysis that follows it will be assumed that $u > 0$. Some remarks about the behavior of h_{ab} near points $X^a(u, v)$ with $u < 0$ will be made towards the end of this section. When $u > 0$ the z -component of the field point satisfies $Z + \frac{1}{2}\pi = u + 4\pi L_0^{-1} \delta z$, and so

$$\chi_{\pm} \equiv v + (u \pm u) + 4\pi L_0^{-1} (\delta t \pm \delta z). \quad (6.7)$$

Furthermore, if $\delta x^a = 0$ the limits of integration in (5.36) are given by $V_0 = v$ and $V_1 = v - W(u)$, where W is the unique root of the equation

$$u - \pi + W = [2 - 2\cos W + (u - \pi)^2]^{1/2}. \quad (6.8)$$

Note that because $V_0 \geq V_1$ it follows that $W \geq 0$, while it is clear from (6.8) that $W = 0$ only at the kink point $u = \pi$. For future reference, it is useful to rearrange (6.8) in the form

$$u - \pi = W^{-1}(1 - \cos W) - \frac{1}{2}W. \quad (6.9)$$

(Incidentally, one way to verify that (6.8) has a unique solution W is to note that the function on the right-hand side of (6.9) is strictly decreasing.)

Implicit differentiation of the defining equation (5.33) for V_k gives

$$\partial_T V_k = H_k^{-1}[T - (k\pi + V_k)]$$

and

$$(\partial_X V_k, \partial_Y V_k, \partial_Z V_k) = H_k^{-1}(\cos V_k - X, \sin V_k - Y, (|k| - \frac{1}{2})\pi - Z), \quad (6.10)$$

with

$$H_k \equiv T - (k\pi + V_k) + X \sin V_k - Y \cos V_k. \quad (6.11)$$

Note that if $\delta x^a = 0$ then $H_0 = u$ and $H_1 = u - \pi + W - \sin W$.

Hence, to linear order in δx^a ,

$$V_0 = v + 4\pi L_0^{-1}(\delta t - \delta z) \quad \text{and} \quad V_1 = v - W + 4\pi L_0^{-1}\delta V \quad (6.12)$$

where

$$\delta V \equiv H_1^{-1}[(u - \pi + W)\delta t + \{\cos(v - W) - \cos v\}\delta x + \{\sin(v - W) - \sin v\}\delta y + (\pi - u)\delta z]. \quad (6.13)$$

Unfortunately, the linear expansion (6.12) for V_0 is not sharp enough to resolve the singular behavior of h_{ab} near the world sheet, as $\chi_- = v + 4\pi L_0^{-1}(\delta t - \delta z)$ and so $\chi_- - V_0 = 0$ at this order.

At quadratic order in δx^a it turns out that

$$V_0 = v + 4\pi L_0^{-1}(\delta t - \delta z) - \frac{1}{2}u^{-1}(4\pi L_0^{-1})^2 Q_+, \quad (6.14)$$

where

$$\begin{aligned} Q_+ &\equiv (\delta x)^2 + (\delta y)^2 + (\delta t - \delta z)(\delta t - \delta z + 2\delta x \sin v - 2\delta y \cos v) \\ &\equiv \bar{\Psi}_{ab}(+1, v) \delta x^a \delta x^b. \end{aligned} \quad (6.15)$$

Since $\bar{\Psi}_{ab}$ is proportional to the orthogonal complement q_{ab} of the projection tensor p_{ab} , the quadratic form Q_+ is zero if and only if δx^a lies in the tangent plane to

the world sheet at $X^a(u, v)$. It should therefore be understood that in evaluating h_{ab} and its derivatives the limit $\delta x^a \rightarrow 0$ is taken only from directions outside the world sheet.

The algebraic computations whose results appear throughout the rest of this section are long and very mechanical, and their details have been relegated to Appendix B. Neglecting contributions that tend to zero in the limit as $\delta x^a \rightarrow 0$, the metric perturbations have the near-field form:

$$(4\mu)^{-1}h_{ab} = \{\ln[(4\pi L_0^{-1})^2 Q_+] - \ln(2u)\}\bar{\Psi}_{ab}(+1, v) + (\gamma_E - \text{Ci } W)\Phi_{ab}(+1, v) \\ + (\text{Si } W)\Omega_{ab}(+1, v) + K(u)\Phi_{ab}(-1, 2u + v) - \Sigma(u)\Omega_{ab}(-1, 2u + v) \\ - (\ln W)\Pi_{ab}(+1) + [\ln(W + 2u - 2\pi) - \ln(2u)]\Pi_{ab}(-1), \quad (6.16)$$

where $\gamma_E \approx 0.5772$ is Euler's constant,

$$\Phi_{ab}(s, v) \equiv \begin{bmatrix} 0 & \sin v & -\cos v & 0 \\ \sin v & 0 & 0 & -s \sin v \\ -\cos v & 0 & 0 & s \cos v \\ 0 & -s \sin v & s \cos v & 0 \end{bmatrix}_{ab}, \quad (6.17)$$

$$\Omega_{ab}(s, v) \equiv \partial_v \Phi_{ab}(s, v) \quad (6.18)$$

and

$$\Pi_{ab}(s) \equiv \begin{bmatrix} 1 & 0 & 0 & -s \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -s & 0 & 0 & 1 \end{bmatrix}_{ab}. \quad (6.19)$$

Note in particular that $\bar{\Psi}_{ab}(s, v) = \Phi_{ab}(s, v) + \Pi_{ab}(s)$. Also, the functions K and Σ are shorthand for

$$K(u) \equiv \text{Ci}(W + 2u - 2\pi) - \text{Ci}(2u) \quad \text{and} \quad \Sigma(u) \equiv \text{Si}(W + 2u - 2\pi) - \text{Si}(2u). \quad (6.20)$$

Similarly, the leading-order contributions to the spacetime derivatives of h_{ab} are:

$$(4\mu)^{-1}(\frac{1}{4\pi}L_0)\partial_t h_{ab} = (\frac{1}{4\pi}L_0)(\partial_t \Lambda)\bar{\Psi}_{ab}(+1, v) + (\delta t - \delta z)(\partial_t \Lambda)\Omega_{ab}(+1, v) \\ + [\Lambda + \gamma_E - \text{Ci } W - \ln(2u)]\Omega_{ab}(+1, v) - (\text{Si } W)\Phi_{ab}(+1, v) \\ + K(u)\Omega_{ab}(-1, 2u + v) + \Sigma(u)\Phi_{ab}(-1, 2u + v) \\ + \frac{\sin W}{u - \pi + W - \sin W}[\frac{1}{W}\bar{\Psi}_{ab}(+1, v - W) \\ - \frac{1}{W + 2u - 2\pi}\bar{\Psi}_{ab}(-1, v - W)], \quad (6.21)$$

$$\begin{aligned}
(4\mu)^{-1}(\frac{1}{4\pi}L_0)\partial_x h_{ab} &= (\frac{1}{4\pi}L_0)(\partial_x \Lambda) \bar{\Psi}_{ab}(+1, v) + (\delta t - \delta z)(\partial_x \Lambda) \Omega_{ab}(+1, v) \\
&+ \frac{\cos(v-W) - \cos v}{u-\pi+W-\sin W} [\frac{1}{W} \bar{\Psi}_{ab}(+1, v-W) \\
&- \frac{1}{W+2u-2\pi} \bar{\Psi}_{ab}(-1, v-W)], \tag{6.22}
\end{aligned}$$

$$\begin{aligned}
(4\mu)^{-1}(\frac{1}{4\pi}L_0)\partial_y h_{ab} &= (\frac{1}{4\pi}L_0)(\partial_y \Lambda) \bar{\Psi}_{ab}(+1, v) + (\delta t - \delta z)(\partial_y \Lambda) \Omega_{ab}(+1, v) \\
&+ \frac{\sin(v-W) - \sin v}{u-\pi+W-\sin W} [\frac{1}{W} \bar{\Psi}_{ab}(+1, v-W) \\
&- \frac{1}{W+2u-2\pi} \bar{\Psi}_{ab}(-1, v-W)] \tag{6.23}
\end{aligned}$$

and

$$\begin{aligned}
(4\mu)^{-1}(\frac{1}{4\pi}L_0)\partial_z h_{ab} &= (\frac{1}{4\pi}L_0)(\partial_z \Lambda) \bar{\Psi}_{ab}(+1, v) + (\delta t - \delta z)(\partial_z \Lambda) \Omega_{ab}(+1, v) \\
&- [\Lambda + \gamma_E - \text{Ci } W - \ln(2u)] \Omega_{ab}(+1, v) + (\text{Si } W) \Phi_{ab}(+1, v) \\
&+ K(u) \Omega_{ab}(-1, 2u+v) + \Sigma(u) \Phi_{ab}(-1, 2u+v) - u^{-1} \bar{\Psi}_{ab}(-1, v) \\
&+ \frac{1}{u-\pi+W-\sin W} [\frac{W-\sin W}{W} \bar{\Psi}_{ab}(+1, v-W) \\
&+ \frac{2u-2\pi+W-\sin W}{W+2u-2\pi} \bar{\Psi}_{ab}(-1, v-W)], \tag{6.24}
\end{aligned}$$

where

$$\Lambda \equiv \ln[(4\pi L_0^{-1})^2 Q_+]. \tag{6.25}$$

The expressions listed here for h_{ab} and its derivatives neglect additional terms of order $\delta x^a \ln[(4\pi L_0^{-1})^2 Q_+]$ or higher, which are zero in the limit $\delta x^a \rightarrow 0$, provided of course that this limit is taken from directions outside the tangent plane to the world sheet at $X^a(u, v)$. However, the expressions also contain terms of order $\ln[(4\pi L_0^{-1})^2 Q_+]$ and $\delta x^a \partial_b \ln[(4\pi L_0^{-1})^2 Q_+]$, which are undefined in the limit as $\delta x^a \rightarrow 0$. The presence of the singular term $4\mu \ln[(4\pi L_0^{-1})^2 Q_+] \bar{\Psi}_{ab}(+1, v)$ in h_{ab} is a reflection of the conical singularity that is expected to mark the location of any zero-thickness cosmic string, as has been explained in more detail in Paper I. Fortunately, none of the singular terms makes a contribution to the first-order equation of motion, as will be seen shortly.

Another important point to note is that equations (6.16) and (6.21)-(6.24) are valid only near points $X^a(u, v)$ with $u \in (0, \pi)$, which lie on one of the two smooth segments of the loop shown in Figure 1. The metric perturbations near points on the second segment, for which $u \in (-\pi, 0)$, can be calculated by making use of the identities $I_+(u - \pi, v) = I_-(u, v)$ and $I_-(u - \pi, v) = I_+(u, v)$ for $u \in (0, \pi)$, and the analogous identities satisfied by the pairs (S_+, S_-) and (C_+, C_-) . In view of equation (5.37) for h_{ab} , it follows that all the components of h_{ab} except h_{tz} , h_{xz}

and h_{yz} are half-range periodic in u near the world sheet (that is, $h_{ab}(u - \pi, v) = h_{ab}(u, v)$), while the components h_{tz} , h_{xz} and h_{yz} are half-range anti-periodic (that is, $h_{ab}(u - \pi, v) = -h_{ab}(u, v)$). This behavior was previously mentioned of the tangential projections $h_{ab}X_{(0),A}^a X_{(0),B}^b$ (which are all half-range periodic in u) in Paper I. Although Paper I deals with a loop that is strictly stationary, the statements it makes about the local behavior of h_{ab} are still applicable here, as the local effects of loop evaporation first appear at order μ^2 in g_{ab} .

6.2. The equations of motion

It is now possible to write down and verify or solve the equations of motion (2.13) and (2.14). Since the only non-zero component of the 2-metric γ^{AB} is $\gamma^{uv} = (\frac{1}{4\pi}L_0)^{-2}$ (to leading order in μ), the zeroth-order equation of motion $q_d^c \gamma^{CD} X_{(0),CD}^d = 0$ reads simply $q_d^c X_{(0),uv}^d = 0$. It is clear by inspection of (4.16) that $X_{(0),uv}^d = 0$, and so (2.13) is automatically satisfied. The second term in the first-order equation of motion (2.14), which is proportional to $\gamma^{CD} X_{(0),CD}^d$, is consequently also zero.

The details of the calculation of the remaining terms in (2.14) can be found in Appendix C. Because all the terms in the first-order equation of motion are projected onto q_d^c , and so lie in the subspace orthogonal to the tangent plane at $X^a(u, v)$, the equation contains only two independent components. In what follows, it proves to be convenient to decompose each term in the directions of the vectors

$$M^a(s, v) \equiv [1, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}} + s \hat{\mathbf{z}}]^a \quad \text{and} \quad N^a(v) \equiv [0, \cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}]^a, \quad (6.26)$$

which span the orthogonal subspace.

The first and third terms in (2.14), which depend on the perturbation δX^a and therefore on the correction functions F , G and A , are:

$$q_d^c \gamma^{CD} \delta X_{(0),CD}^d = 2\mu(\frac{1}{4\pi}L_0)^{-1}(F_{uv} + sA_{uv} + G_u - \kappa) M^c + 2\mu(\frac{1}{4\pi}L_0)^{-1}G_{uv} N^c \quad (6.27)$$

and

$$-2q_d^c \gamma^{AC} \gamma^{BD} \eta_{ab} X_{(0),A}^a \delta X_{(0),B}^b X_{(0),CD}^d = 2\mu(\frac{1}{4\pi}L_0)^{-1}[\kappa s(|u| - \frac{1}{2}\pi) - (F_u + sA_u)]N^c. \quad (6.28)$$

The remaining terms depend on h_{ab} and its derivatives, which were evaluated earlier in the case $u \in (0, \pi)$ only. For positive values of u (or equivalently, for

$s = +1$) it turns out that

$$-q_d^c \gamma^{AC} \gamma^{BD} h_{ab} X_{(0),A}^a X_{(0),B}^b X_{(0),CD}^d = 16\mu(\frac{1}{4\pi}L_0)^{-1}[\ln(W + 2u - 2\pi) - \ln(2u)]N^c, \quad (6.29)$$

$$\begin{aligned} -q^{cd} \gamma^{AB} X_{(0),A}^a X_{(0),B}^b h_{bd,a} &= 16\mu(\frac{1}{4\pi}L_0)^{-1} \Sigma(u) [-(2 \cos 2u)M^c + (\sin 2u)N^c] \\ &+ 16\mu(\frac{1}{4\pi}L_0)^{-1} K(u) [(2 \sin 2u)M^c + (\cos 2u)N^c] \\ &+ 4\mu(\frac{1}{4\pi}L_0)^{-1} \frac{1}{u-\pi+W-\sin W} [(1 - \cos W)M^c - (\sin W)N^c] \\ &+ 4\mu(\frac{1}{4\pi}L_0)^{-1} \frac{W+2u-2\pi-2\sin W}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} \\ &\times [3(1 - \cos W)M^c - (\sin W)N^c] \end{aligned} \quad (6.30)$$

and

$$\begin{aligned} \frac{1}{2}q^{cd} \gamma^{AB} X_{(0),A}^a X_{(0),B}^b h_{ab,d} &= -16\mu(\frac{1}{4\pi}L_0)^{-1} [K(u) \sin 2u - \Sigma(u) \cos 2u]M^c \\ &- 8\mu(\frac{1}{4\pi}L_0)^{-1} \frac{W+2u-2\pi-\sin W}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi} M^c \\ &- 8\mu(\frac{1}{4\pi}L_0)^{-1} \frac{1-\cos W}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi} N^c. \end{aligned} \quad (6.31)$$

Now, it was shown in Paper I that $h_{ab}X_{(0),A}^a X_{(0),B}^b$ is half-range periodic in u , and it is easily seen that taking the inner product with $q_d^c \gamma^{AC} \gamma^{BD} X_{(0),CD}^d$ preserves this symmetry. Hence, the functional dependence of the term on the left of (6.29) for $u \in (-\pi, 0)$ is the half-range periodic extension of the term on the right. Similarly, the sum of the terms in (6.30) and (6.31) – which is to say, the term on the right-hand side of the first-order equation of motion (2.14) – is just $2\alpha^a$, where α^a is the acceleration vector introduced in Paper I. It was demonstrated in Paper I that the t -, x - and y -components of α^a are half-range periodic in u , while the z -component is half-range anti-periodic, and this is also true of the vectors $M^c(s, v)$ and $N^c(v)$. Hence, the first-order equation of motion requires the sum of (6.27) and (6.28) to have the same symmetries, and since the term $\kappa s(|u| - \frac{1}{2}\pi)$ in (6.28) is half-range periodic this means that $F + sA$ and G must be half-range periodic as well.

These properties can also be predicted from the geometry of the problem. The two segments of the evaporating loop should share the same geometry, so the correction functions F , G and A are expected to have the same symmetries as the corresponding components of the original ACO loop. That is, F and G should be half-range periodic in u as the t and x - y components of the ACO loop are, while A should be half-range anti-periodic as the z -component $|u| - \frac{1}{2}\pi$ is.

On setting $s = +1$ in the first two terms (6.27) and (6.28), the equation of motion for $u \in (0, \pi)$ reduces to two partial differential equations:

$$F_{uv} + A_{uv} + G_u - \kappa = 8[K(u) \sin 2u - \Sigma(u) \cos 2u] + 4 \frac{1 - \cos W}{u - \pi + W - \sin W} \frac{W + 2u - 2\pi - 2 \sin W}{W + 2u - 2\pi} \quad (6.32)$$

and

$$\begin{aligned} & G_{uv} + \kappa(u - \tfrac{1}{2}\pi) - (F_u + A_u) + 8[\ln(W + 2u - 2\pi) - \ln(2u)] \\ &= 8[K(u) \cos 2u + \Sigma(u) \sin 2u] - 4 \frac{1}{u - \pi + W - \sin W} (\sin W - 2 \frac{1 - \cos W}{W + 2u - 2\pi} \cos W) \end{aligned} \quad (6.33)$$

which appear as the coefficients of M^c and N^c respectively.

At a schematic level, these equations read:

$$F_{uv} + A_{uv} + G_u = P_1(u) \quad \text{and} \quad G_{uv} - F_u - A_u = P_2(u), \quad (6.34)$$

where the functions P_1 and P_2 represent all the remaining terms. Taking the v derivative of each equation in turn and adding or subtracting the other leads to two forced simple harmonic equations:

$$G_{uvv} + G_u = P_1(u) \quad \text{and} \quad (F_u + A_u)_{vv} + F_u + A_u = -P_2(u), \quad (6.35)$$

whose general solutions are:

$$G = c_1(v) + a(u) \cos v + b(u) \sin v + \int P_1(u) du \quad (6.36)$$

and

$$F + A = c_2(v) + b(u) \cos v - a(u) \sin v - \int P_2(u) du, \quad (6.37)$$

where c_1 and c_2 are arbitrary continuous 2π -periodic functions of v , and a and b are the restrictions to $(0, \pi)$ of arbitrary half-range periodic C^1 functions of u .

The physical significance of the arbitrary homogeneous contributions to (6.36) and (6.37) can be understood by referring to the intrinsic 2-metric γ_{AB} . The components of γ_{AB} are expanded to order μ in Appendix D.1, where it is shown that γ_{AB} takes on the self-similar form (4.22) at this order if and only if

$$F_{uv} + A_{uv} = 0, \quad F_{uv} + F_{vv} + A_{vv} = 0 \quad \text{and} \quad G_v + F_{vv} = 0. \quad (6.38)$$

The first of these constraints reduces the first equation in (6.34) to $G_u = P_1(u)$, and so from (6.36) a and b are both zero. The trigonometric functions of v in

(6.36) and (6.37) therefore represent free vibrations of the loop with amplitudes of order μ , which break the self-similarity of the evaporating loop and, if allowed to remain, would contribute to the metric tensor g_{ab} at order μ^2 .

The second of the constraints (6.38) now reduces to $F_{uv} = -c_2''(v)$, and since F is by assumption continuous and half-range periodic in u it follows that $c_2' = 0$ and therefore that c_2 is a constant (and can be absorbed into $\int P_2(u) du$). Consequently $F_{uv} = 0$, and F has the generic form $F_1(u) + F_2(v)$ for some choice of continuous and suitably periodic functions F_1 and F_2 . Furthermore, the third constraint now becomes $G_v = -F_2''$, and so from (6.36) $c_1(v) = -F_2'$.

The possibility of further constraining F , G and A arises from a consideration of transformations of the spacetime coordinates $x^a = [t, x, y, z]^a$ of the form $x^a \rightarrow \hat{x}^a$ where

$$\hat{x}^a = x^a + \mu y^a(x^b). \quad (6.39)$$

The only transformations of this type that are admissible are those that preserve the form of the metric tensor $g_{ab} = \eta_{ab} + \mu h_{ab}$ at order μ . Since

$$\hat{g}_{ab} = \eta_{ab} + \mu h_{ab} + \mu \eta_{ca} \partial_b y^c + \mu \eta_{cb} \partial_a y^c \quad (6.40)$$

at linear order in μ , admissible transformations satisfy the condition $\eta_{c(a} \partial_{b)} y^c = 0$ and so are generators of the Poincaré transformations (that is, combinations of boosts, rotations and translations).

Under such a transformation, the position vector X^a of the loop is replaced by

$$\hat{X}^a = X_{(0)}^a + \delta X^a + \mu Y^a(u, v) \quad (6.41)$$

where $X_{(0)}^a$ and δX^a are given by (C.1) and (C.3) respectively, and

$$Y^a(u, v) \equiv \frac{1}{4\pi} L_0 [-\delta F(u, v), \delta G(u, v) \{\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}\} + \delta A(u, v) \hat{\mathbf{z}}]^a \quad (6.42)$$

for some choice of functions δF , δG and δA .⁶

It is only possible to impose the admissibility condition $\eta_{c(a} \partial_{b)} y^c = 0$ on Y^a if the derivatives are taken tangent to the worldsheet of the loop, and so Y^a must satisfy the equations

$$X_{(0),u} \cdot Y_{,u} = 0, \quad X_{(0),v} \cdot Y_{,v} = 0 \quad \text{and} \quad X_{(0),u} \cdot Y_{,v} + X_{(0),v} \cdot Y_{,u} = 0. \quad (6.43)$$

⁶Note that the x - and y -components of Y^a have the forms shown so as to preserve the isotropic gauge condition discussed in Section 4, following equation (4.20).

In terms of δF , δG and δA these equations read, for $u \in (0, \pi)$,

$$\delta F_u + \delta A_u = 0 \quad \delta G + \delta F_v = 0 \quad \text{and} \quad \delta F_v + \delta A_v + \delta F_u = 0. \quad (6.44)$$

The choice $\delta F = -F_2(v)$, $\delta G = F'_2(v)$ and $\delta A = F_2(v)$ satisfies all three equations, and specifies a spacetime transformation which reduces F and A to functions of u only, and sets $c_1 = 0$ in (6.36).

In a suitable set of spacetime coordinates, therefore, the correction functions become

$$G = \int P_1(u) du \quad \text{and} \quad F + A = - \int P_2(u) du, \quad (6.45)$$

with F a continuous and half-range periodic function of u alone. In fact, F is effectively an arbitrary function of u , as it is always possible to define a transformation $u \rightarrow u' = u + \mu U(u)$ of the gauge coordinate u , where U is continuous and half-range periodic but otherwise arbitrary, under which $X^a \rightarrow X^a + \mu U X^a_{(0),u}$ and so $F \rightarrow F - U$ and $A \rightarrow A + U$ for $u \in (0, \pi)$, but G remains unchanged.

6.3. Solving for F , G and A

The equation (6.9) specifying $u - \pi$ as a function of W can be used to eliminate u from the terms in (6.32) and (6.33) which mix both u and W . The resulting expressions for P_1 and P_2 read:

$$P_1(u) = \kappa + 8[K(u) \sin 2u - \Sigma(u) \cos 2u] + 4(1 - \cos W - W \sin W)/J(W) \quad (6.46)$$

and

$$\begin{aligned} P_2(u) = & -\kappa(u - \tfrac{1}{2}\pi) - 8[\ln\{2W^{-1}(1 - \cos W)\} - \ln(2u)] \\ & + 8[K(u) \cos 2u + \Sigma(u) \sin 2u] - 4(\sin W - W \cos W)/J(W), \end{aligned} \quad (6.47)$$

where

$$J(W) \equiv W^{-1}(1 - \cos W) + \tfrac{1}{2}W - \sin W \quad (6.48)$$

and now

$$K(u) = \text{Ci}\{2W^{-1}(1 - \cos W)\} - \text{Ci}(2u) \quad \text{and} \quad \Sigma(u) = \text{Si}\{2W^{-1}(1 - \cos W)\} - \text{Si}(2u). \quad (6.49)$$

Furthermore, the derivative of (6.9) reads $du = -W^{-1}J(W)dW$, and so

$$\begin{aligned} G &= \kappa u + 8 \int [K(u) \sin 2u - \Sigma(u) \cos 2u] du - 4 \int W^{-1}(1 - \cos W - W \sin W) dW \\ &= \kappa u - 4[K(u) \cos 2u + \Sigma(u) \sin 2u] + 4 \ln(1 - \cos W) - 4 \ln(uW) + C_1 \end{aligned} \quad (6.50)$$

and

$$\begin{aligned} F + A &= \frac{1}{2}\kappa u(u - \pi) + 8 \int [\ln\{2W^{-1}(1 - \cos W)\} - \ln(2u)] du \\ &\quad - 8 \int [K(u) \cos 2u + \Sigma(u) \sin 2u] du - 4 \int W^{-1}(\sin W - W \cos W) dW \\ &= \frac{1}{2}\kappa u(u - \pi) - 8W^{-1}[1 + \ln W - \ln(1 - \cos W)](1 - \cos W) - 4W(2 - \ln W) \\ &\quad - 8(u \ln u - u) - 4[K(u) \sin 2u - \Sigma(u) \cos 2u] - 4 \int_{2\pi}^W \ln(1 - \cos \omega) d\omega + C_2. \end{aligned} \quad (6.51)$$

Here, the integral $\int \ln(1 - \cos \omega) d\omega$ unfortunately cannot be expressed in closed form as a combination of standard functions. (The lower bound in the integral has been set to $\omega = 2\pi$ because $W \rightarrow 2\pi^-$ as $u \rightarrow 0^+$.) Equations (6.50) and (6.51) can of course be verified directly by differentiation.. However, in tracing out the original calculations – that is, in integrating P_1 and P_2 – it proves to be most efficient to integrate the terms $K(u) \sin 2u - \Sigma(u) \cos 2u$ in (6.50) and $\ln(1 - \cos W)$ and $\Sigma(u) \sin 2u + K(u) \cos 2u$ in (6.51) by parts with respect to u first, then integrate the remainders with respect to W .

Turning to the initial conditions satisfied by G and $F + A$, it should be recalled that G is continuous and half-range periodic in u and so $G(0) = G(\pi)$. Similarly, F is half-range periodic in u and so $F(0) = F(\pi)$. By contrast, A is continuous and half-range *anti*-periodic in u , and therefore $A(0) = -A(\pi)$. (Note that the combination $F + sA$, which appears in the leading terms (6.27) and (6.28) in the first-order equation of motion, is half-range periodic for $u \in [-\pi, \pi]$ as required. However, it is not necessarily continuous at $u = 0$.)

Now, the $u \rightarrow 0^+$ lower limit of equation (6.51) reads:

$$F(0) + A(0) = -16\pi + 8\pi \ln 2\pi + C_2 \quad (6.52)$$

while the $u \rightarrow \pi^-$ upper limit reads:

$$F(\pi) + A(\pi) = 8\pi - 8\pi \ln(2\pi) - 4 \text{Si}(2\pi) + C_2. \quad (6.53)$$

(Note in particular that $\int_{2\pi}^0 \ln(1 - \cos \omega) d\omega = 2\pi \ln 2$ at the upper limit.)

Subtracting (6.53) from (6.52) therefore gives:

$$2A(0) \equiv A(0) - A(\pi) = -24\pi + 16\pi \ln 2\pi + 4 \operatorname{Si}(2\pi) \quad (6.54)$$

and so

$$A(0) = -12\pi + 8\pi \ln 2\pi + 2 \operatorname{Si}(2\pi) \approx 11.328. \quad (6.55)$$

Similarly, addition of (6.52) and (6.53) gives:

$$2F(0) \equiv F(0) + F(\pi) = -8\pi - 4 \operatorname{Si}(2\pi) + 2C_2 \quad (6.56)$$

and so

$$F(0) = -4\pi - 2 \operatorname{Si}(2\pi) + C_2 \approx -15.403 + C_2. \quad (6.57)$$

The constant of integration C_2 in (6.57) represents a vestige of coordinate freedom in the specification of the correction function F that, in view of (6.55) is not present in the function A . In fact, different values of C_2 correspond to the same physical configuration of the evaporating loop, as the value of C_2 can be changed at will by making a suitable Poincaré transformation (6.41). This follows from the fact that the choice $\delta G = \delta A = 0$ and δF any arbitrary constant (representing a translation in the coordinate t at order μ) automatically satisfies the Poincaré constraints (6.44). One natural gauge choice for F is to set $F(0) = 0$, in which case $C_2 = 4\pi + 2 \operatorname{Si}(2\pi)$.

Consider now equation (6.50) for G . Taking the lower limit $u \rightarrow 0^+$ (or equivalently $W \rightarrow 2\pi^-$) in this equation gives:

$$G(0) = C_1, \quad (6.58)$$

while taking the upper limit $u \rightarrow \pi^-$ (equivalently $W \rightarrow 0^+$) gives:

$$G(\pi) = \pi\kappa + 4 \operatorname{Ci}(2\pi) - 4 \ln 2\pi - 4\gamma_E + C_1. \quad (6.59)$$

So it follows that $G(0) = G(\pi)$ if and only if

$$\kappa = [4 \ln 2\pi + 4\gamma_E - 4 \operatorname{Ci}(2\pi)]/\pi \approx 3.1037. \quad (6.60)$$

That is, iff $\kappa = \gamma^0/(4\pi)$ with $\gamma^0 \approx 39.0025$ the radiative efficiency of the ACO loop, as expected.

Unlike C_2 , the integration constant C_1 in (6.50) does have an invariant physical significance, and it cannot be arbitrarily changed by making a Poincaré or gauge

transformation without affecting the form of F . In fact, different values of C_1 correspond to different values for the length of the evaporating loop as measured along curves of constant scale factor $e^{-\kappa\mu(u+v)}$, as is explained in more detail in Appendix D.2. If the length L_{csf} of the loop along such a curve is assumed to be independent of μ (with $L_{\text{csf}} = \frac{1}{\sqrt{2}}L_0$ along the curve $u + v = 0$) then

$$C_1 = -28 - 6\gamma_E + 10 \ln 2\pi + 6 \text{Ci}(2\pi) + 4\pi^{-1} \text{Si}(2\pi) + 4\pi^{-1} \text{Si}(4\pi) \approx -9.5143. \quad (6.61)$$

With C_1 assigned this value, the correction function G , which is the half-range periodic extension of the function on the right-hand side of (6.50), is plotted for $u \in [-\pi, \pi]$ in Figure 5a. The half-range periodic extension to $[-\pi, \pi]$ of the function $F + A$ defined by equation (6.51) with $C_2 = 4\pi + 2 \text{Si}(2\pi)$ is plotted against u in Figure 5b. As mentioned at the end of Section 6.2, the functions F and A are not separately uniquely determined, as different choices of F correspond to different choices of the gauge coordinate u . However, the requirement that $A(0) = -A(\pi) = \lambda$ (with $\lambda \equiv -12\pi + 8\pi \ln 2\pi + 2 \text{Si}(2\pi)$) does place a restriction on the choice of F . One natural gauge choice is to set A proportional to the z -component $|u| - \frac{1}{2}\pi$ of the unperturbed position function. Then

$$A(u) = -\frac{2\lambda}{\pi}(|u| - \frac{1}{2}\pi) \quad (6.62)$$

is continuous and half-range anti-periodic on $[-\pi, \pi]$ with $A(0) = \lambda$, while from (6.51) F is the half-range periodic extension to $[-\pi, \pi]$ of

$$\begin{aligned} F(u) = & \frac{1}{2}\kappa u(u - \pi) + \frac{2\lambda}{\pi}(u - \frac{1}{2}\pi) - 8W^{-1}[1 + \ln W - \ln(1 - \cos W)](1 - \cos W) \\ & - 4W(2 - \ln W) - 8(u \ln u - u) - 4[K(u) \sin 2u - \Sigma(u) \cos 2u] \\ & - 4 \int_{2\pi}^W \ln(1 - \cos \omega) d\omega + 4\pi + 2 \text{Si}(2\pi). \end{aligned} \quad (6.63)$$

The correction functions A and F in this gauge are plotted against u in Figures 6a and 6b respectively.

To summarize, the position function (4.20) with F , G and A suitable extensions of (6.63), (6.50) and (6.62) is a solution of the first-order string equation of motion (2.14) that reduces to the ACO solution in the limit as $\mu \rightarrow 0$. Furthermore, if the integration constant C_1 is chosen to take the value (6.61) then the solution is the unique self-similar loop trajectory (up to gauge and Poincaré transformations) with the property that the length of the loop retains its ACO value $\frac{1}{\sqrt{2}}L_0$ along the curve $u + v = 0$ (which to leading order in μ corresponds to the spacelike slice $t = 0$).

7. Matching the Solution to the Minkowski Vacuum

By prescription, the evaporating ACO loop radiates away the last of its energy at $t = t_L$, the moment when it shrinks to a point at the origin. It is therefore to be expected that the metric induced by the loop reduces to the Minkowski vacuum at all points in the chronological future of the evaporation point $x^a = [t_L, \mathbf{0}]^a$. If the evaporation point is denoted by p_L , the boundary of the chronological future $I^+(p_L)$ is the future light cone F_L of p_L , plus p_L itself. The weak-field metric will match smoothly onto Minkowski spacetime in $I^+(p_L)$ if an appropriate set of junction conditions is satisfied on F_L . I will show explicitly that these junction conditions are indeed satisfied later in this section. (Note, however, that the weak-field metric remains singular in every neighborhood of p_L , and cannot be matched onto Minkowski spacetime at this one point.)

7.1. The surface $t - t_L = |\mathbf{x}|$ is null to order μ

Before proceeding, it is important to first establish that the surface F_L is, to order μ , just the surface $t - t_L = |\mathbf{x}|$, which is of course null with respect to the unperturbed metric tensor η_{ab} . To show that the surface $t - t_L = |\mathbf{x}|$ is also null with respect to the first-order metric tensor $\eta_{ab} + h_{ab}$, it is sufficient to show that $h_{ab}(t, \mathbf{x}) \rightarrow 0$ as the field point $x^a = [t, \mathbf{x}]^a$ approaches the surface.

Suppose then that ε is a number in $(0, 1)$, and consider the set of field points $[t, \mathbf{x}]$ lying to the past of the surface $t - t_L = |\mathbf{x}|$ and satisfying the conditions

$$|\mathbf{x}| - t + t_L = \varepsilon t_L \quad \text{and} \quad [(t - t_L)^2 + |\mathbf{x}|^2]^{1/2} > \varepsilon^{1/2} t_L. \quad (7.1)$$

The first condition implies that ε is a dimensionless measure of the distance from the point $[t, \mathbf{x}]$ to the surface $t - t_L = |\mathbf{x}|$ in the background Minkowski manifold. In fact, it is easily seen that the *Euclidean* distance from the point to the surface is just $\frac{1}{\sqrt{2}}\varepsilon t_L$. The second condition ensures that the field point avoids a certain neighborhood of the evaporation point $[t_L, \mathbf{0}]$. (The scaling factor $\varepsilon^{1/2}$ is chosen here because the exclusion of a neighborhood about $[t_L, \mathbf{0}]$ with the smaller radius εt_L does not lead to a useful bound on the components of h_{ab} , as will be seen shortly.) It is evident that as $\varepsilon \rightarrow 0$ the points satisfying (7.1) include all points in some timelike past neighborhood of every point on the surface $t - t_L = |\mathbf{x}|$ except the evaporation point.

Eliminating t from the conditions (7.1) yields one non-trivial bound on $|\mathbf{x}|$ if $\varepsilon < 1$, namely

$$|\mathbf{x}| > \frac{1}{2}[\varepsilon + (2\varepsilon - \varepsilon^2)^{1/2}] t_L, \quad (7.2)$$

which in turn implies that $|\mathbf{x}| > \frac{1}{\sqrt{2}}\varepsilon^{1/2}t_L$ (as can readily be verified by dividing the expression on the right of (7.2) by $\varepsilon^{1/2}t_L$ and showing that the minimum value of the resulting function is $1/\sqrt{2}$).

Now, the corresponding image point $[\bar{t}, \bar{\mathbf{x}}]$ on the future light cone F_0 of the origin has coordinates

$$\bar{t} = e^{\kappa\mu\psi}|\mathbf{x}| \quad \text{and} \quad \bar{\mathbf{x}} = e^{\kappa\mu\psi}R(-\psi)\mathbf{x} \quad (7.3)$$

where

$$e^{\kappa\mu\psi} \equiv [(|\mathbf{x}| - t + t_L)/t_L]^{-1} = \varepsilon^{-1} \quad (7.4)$$

and R is the rotation matrix defined in (5.14). So in particular

$$\bar{t} = \varepsilon^{-1}|\mathbf{x}| > \frac{1}{\sqrt{2}}\varepsilon^{-1/2}t_L, \quad (7.5)$$

while of course $|\bar{\mathbf{x}}| = \bar{t}$.

Note here that $\bar{t} \rightarrow \infty$ as $\varepsilon \rightarrow 0$. Also, according to (5.22),

$$h_{ab}(t, \mathbf{x}) = R_a^c(\psi) h_{cd}(\bar{t}, \bar{\mathbf{x}}) R_b^d(\psi) \quad (7.6)$$

where the components of the rotation operators R_b^a are all bounded functions of ψ . Hence, to prove that $h_{ab}(t, \mathbf{x}) \rightarrow 0$ as $\varepsilon \rightarrow 0$ in the conditions (7.1), it is sufficient to show that $|h_{ab}(\bar{t}, \bar{\mathbf{x}})|$ is bounded above on F_0 by a function of $\bar{t}$ which goes to zero (uniformly with respect to $\bar{\mathbf{x}}$) as $\bar{t} \rightarrow \infty$. It should also be clear that a proof of the last statement would not be sufficient if the factor $\varepsilon^{1/2}$ in the second condition in (7.1) were replaced by ε or a multiple of ε . For then (7.2) would be replaced by an inequality of the form $|\mathbf{x}| > c\varepsilon$ for some constant c , and the image point $[\bar{t}, \bar{\mathbf{x}}]$ would be placed under the much weaker constraint $\bar{t} > c$, which does not guarantee that $h_{ab}(\bar{t}, \bar{\mathbf{x}}) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Now, given that the components of $\bar{\Psi}_{ab}$ in the equation (5.36) for $h_{ab}(\bar{t}, \bar{\mathbf{x}})$ all satisfy $|\bar{\Psi}_{ab}(s, v)| \leq 1$, it follows that

$$\begin{aligned} |h_{ab}(\bar{t}, \bar{\mathbf{x}})| &\leq 4\mu \int_{V_0}^{V_1+2\pi} [T + (Z + \tfrac{1}{2}\pi) - v]^{-1} dv + 4\mu \int_{V_1}^{V_0} [T - (Z + \tfrac{1}{2}\pi) - v]^{-1} dv \\ &\equiv 4\mu|I_+| + 4\mu|I_-|, \end{aligned} \quad (7.7)$$

where $[T, X, Y, Z] = 4\pi L_0^{-1}[\bar{t}, \bar{\mathbf{x}}]$ as before. So $|h_{ab}(\bar{t}, \bar{\mathbf{x}})|$ will be bounded above by a function which goes uniformly to zero as $\bar{t} \rightarrow \infty$ if the same is true, separately, of $|I_+|$ and $|I_-|$.

Consider then

$$|I_-| = \int_{V_1}^{V_0} [E(1, v)]^{-1} dv \quad \text{where} \quad E(s, v) = T - s(Z + \tfrac{1}{2}\pi) - v. \quad (7.8)$$

Since E is positive semi-definite (see Appendix A.2), there are two possible cases that need to be considered: either (i) $E(1, v) = 0$ for some $v \in [V_1, V_0]$, or (ii) $E(1, v) > 0$ for all $v \in [V_1, V_0]$.

Case (i) has been treated in some detail in Appendix A.2, where it is shown that $E(1, v) = 0$ for some v only if the point (X, Y) lies on the unit circle (so that $T = (1 + Z^2)^{1/2}$) and $Z > -\frac{1}{2}\pi$. Furthermore, if $Z > \frac{1}{2}\pi$ (and so $T > (1 + \frac{1}{4}\pi^2)^{1/2} \approx 1.8621$, which will henceforth be assumed) then the values of V_0 and V_1 coincide, and according to (A.9)

$$|I_-| = \ln[1 + \pi/(Z - \tfrac{1}{2}\pi)]. \quad (7.9)$$

In view of the inequality $\ln(1 + x) \leq x$ it follows that

$$|I_-| \leq \pi/(Z - \tfrac{1}{2}\pi) \equiv \pi/[(T^2 - 1)^{1/2} - \tfrac{1}{2}\pi] \quad (7.10)$$

where the function of T appearing on the right is bounded above by $8\pi/T$ if $T \geq \frac{1}{63}(4\pi + 8\sqrt{16\pi^2 + 63}) \approx 2.0869$. (The significance of the factor 8π will become apparent shortly.) So, in this case,

$$|I_-| \leq 8\pi/T \quad \text{if} \quad T \gtrsim 2.0869. \quad (7.11)$$

In case (ii), $E(1, v) > 0$ for all $v \in [V_1, V_0]$. Since E is a linear decreasing function of v , there exists a number V_* in $[V_1, V_0]$ with the property that $E(1, v) > \frac{1}{2}T$ for all $v \in (V_1, V_*)$ and $0 < E(1, v) < \frac{1}{2}T$ for all $v \in (V_*, V_0)$.⁷ An immediate consequence is that

$$|I_-| = \int_{V_1}^{V_*} [E(1, v)]^{-1} dv + \int_{V_*}^{V_0} [E(1, v)]^{-1} dv \quad (7.12)$$

where

$$\int_{V_1}^{V_*} [E(1, v)]^{-1} dv < 2(V_* - V_1)/T \leq 4\pi/T, \quad (7.13)$$

⁷The value of V_* will of course depend on $[T, X, Y, Z]$. Also, V_* may coincide with either V_0 or V_1 , in which case one of the two domains specified for v will be empty.

as $V_* - V_1 \leq V_0 - V_1 \leq 2\pi$.

The integral over the remaining segment (V_*, V_0) can be bounded above by using equation (5.34) to replace v with u as the integration variable, so that

$$\int_{V_*}^{V_0} [E(1, v)]^{-1} dv = \int_0^{u(V_*)} [H(v)]^{-1} du \quad (7.14)$$

where $H(v) = T - (u + v) + X \sin v - Y \cos v$ is a second positive semi-definite function, examined in some detail in Appendix A.3. In particular, it is evident from the first line of (A.10) that

$$H(v) \geq |Z - (|u| - \tfrac{1}{2}\pi)| \geq Z - \tfrac{1}{2}\pi \quad (7.15)$$

if u is in $[0, \pi]$ and $Z > \frac{1}{2}\pi$. Also, the condition $E(1, v) < \frac{1}{2}T$ corresponds to the inequality

$$Z > \tfrac{1}{2}T - \tfrac{1}{2}\pi - v \geq \tfrac{1}{2}T - \tfrac{3}{2}\pi - (1 + \tfrac{1}{4}\pi^2)^{1/2} \quad (7.16)$$

where v here has been replaced by its maximum possible value, $\pi + (1 + \frac{1}{4}\pi^2)^{1/2}$ [see Appendix A.1]. In particular, the condition $Z > \frac{1}{2}\pi$ will be satisfied if $T \gtrsim 16.2906$. Hence, if the latter condition is assumed,

$$\int_0^{u(V_*)} [H(v)]^{-1} du < u(V_*) [\tfrac{1}{2}T - 2\pi - (1 + \tfrac{1}{4}\pi^2)^{1/2}]^{-1} \quad (7.17)$$

where the maximum possible value of $u(V_*)$ is π . The function of T appearing on the right of (7.17) is therefore bounded above by $4\pi/T$ if $T \geq 8\pi + 2\sqrt{4 + \pi^2} \approx 32.5811$.

In summary, the inequalities (7.13) and (A.10) together imply that

$$|I_-| \leq 8\pi/T \quad \text{if} \quad T \gtrsim 32.5811 \quad (7.18)$$

in case (ii), and so in view of (7.11) $|I_-|$ is bounded above by a function that goes to zero uniformly as $T \rightarrow \infty$, as required. Note that the factor of 8π in the bounding function arises as a consequence of choosing V_* to separate the domains $E \leq \frac{1}{2}T$. The factor $\frac{1}{2}$ is to some extent arbitrary here, as any factor less than 1 will induce similar bounds on $|I_-|$. However, a factor 1 or greater is not viable, as it generates no useful bound on Z , and hence no useful bound on the integral (7.14).

The proof that the second contribution $|I_+|$ to $|h_{ab}(\bar{t}, \bar{\mathbf{x}})|$ is also uniformly bounded above by $8\pi/T$ for sufficiently large values of T is formally identical to

the proof just given, provided that $|I_-|$ is everywhere replaced with $|I_+|$, $E(1, v)$ with $E(-1, v)$, Z with $-Z$, V_0 with $V_1 + 2\pi$ and V_1 with V_0 . Also, the limit $u = 0$ in the integral on the right of (7.14) should there and below be replaced with $u = -\pi$, and the bound $u(V_*) \leq \pi$ with $u(V_*) \leq 0$.

In conclusion, therefore, $|h_{ab}(\bar{t}, \bar{\mathbf{x}})| \leq 64\pi\mu/T \equiv 16L_0/\bar{t}$ for sufficiently large values of $\bar{t}$, and so $h_{ab}(t, \mathbf{x}) \rightarrow 0$ as $\varepsilon \rightarrow 0$ in the conditions (7.1). It follows that the future light cone F_L of the final evaporation point is, to order μ , just the surface $t - t_L = |\mathbf{x}|$.

7.2. Checking the junction conditions

The junction conditions that apply when matching two spacetime regions across a null hypersurface $\mathcal{N}$ were first usefully formulated by Barrabès and Israel [20] and have been handily summarized by Poisson in [21]. The first step in the Barrabès-Israel method is to parametrize the null hypersurface in the form $x^a = y^a(r, \theta^1, \theta^2)$, where r is an arbitrary coordinate along the generators of the hypersurface, and each generator is labeled by the two remaining coordinates θ^A (for $A = 1, 2$). This parametrization can in turn be used to construct an orthonull basis $\{k^a, e_1^a, e_2^a, N^a\}$ at each point on $\mathcal{N}$, with

$$k^a = \partial y^a / \partial r, \quad e_A^a = \partial y^a / \partial \theta^A \quad (7.19)$$

and the remaining (null) basis field N^a uniquely determined by the constraints

$$g_{ab}N^a k^b = 1 \quad \text{and} \quad g_{ab}N^a e_A^b = 0. \quad (7.20)$$

The hypersurface then has an associated mass density ρ , current density j^A and isotropic pressure p given by

$$\rho = -\frac{1}{8\pi}\sigma^{AB}[C_{AB}], \quad j^A = \frac{1}{8\pi}\sigma^{AB}[C_{3B}] \quad \text{and} \quad p = -\frac{1}{8\pi}[C_{33}], \quad (7.21)$$

where $\sigma_{AB} = g_{ab}e_A^a e_B^b$ and $C_{\alpha\beta} = e_\alpha^a e_\beta^b \nabla_b N_a$, with α, β running from 1 to 3 and $e_3^a \equiv k^a$. Also, $[C_{\alpha\beta}]$ denotes $C_{\alpha\beta}(\mathcal{N}^+) - C_{\alpha\beta}(\mathcal{N}^-)$ where $\mathcal{N}^-$ and $\mathcal{N}^+$ are the (timelike) past- and future-facing sides of $\mathcal{N}$ respectively. (Note that $C_{\alpha\beta}$ typically has different values on the two sides of $\mathcal{N}$ because the limiting values of the covariant derivatives $\nabla_b N_a$ are different, even though the metric tensor g_{ab} is assumed here to be continuous across $\mathcal{N}$.) The two spacetime regions can be matched across $\mathcal{N}$ without invoking an extraneous source of stress-energy on $\mathcal{N}$ if ρ , j^A and p are all zero.

In the case of the evaporating ACO loop, $\mathcal{N}$ is the null hypersurface F_L defined by $t - t_L = |\mathbf{x}|$, and $g_{ab} = \eta_{ab}$ there. An obvious choice of parametrization is

$$y^a(r, \theta^1, \theta^2) = [t_L, \mathbf{0}]^a + [r, r \sin \theta^1 \cos \theta^2, r \sin \theta^1 \sin \theta^2, r \cos \theta^1]^a. \quad (7.22)$$

The corresponding orthonull basis fields are:

$$k^a = [1, \sin \theta^1 \cos \theta^2, \sin \theta^1 \sin \theta^2, \cos \theta^1]^a, \quad (7.23)$$

$$e_1^a = [0, r \cos \theta^1 \cos \theta^2, r \cos \theta^1 \sin \theta^2, -r \sin \theta^1]^a, \quad (7.24)$$

$$e_2^a = [0, -r \sin \theta^1 \sin \theta^2, r \sin \theta^1 \cos \theta^2, 0]^a \quad (7.25)$$

and

$$N^a = \frac{1}{2}[1, -\sin \theta^1 \cos \theta^2, -\sin \theta^1 \sin \theta^2, -\cos \theta^1]^a. \quad (7.26)$$

In particular, the line element $\sigma_{AB} d\theta^A d\theta^B$ intrinsic to the hypersurface is just that of a spatial 2-sphere:

$$\sigma_{AB} = -r^2 \text{diag}(1, \sin^2 \theta^1) \quad \text{and} \quad \sigma^{AB} = -r^{-2} \text{diag}(1, \csc^2 \theta^1). \quad (7.27)$$

To linear order in the derivatives of h_{ab} , the transverse curvature of F_L is

$$C_{\alpha\beta} = e_\alpha^a e_\beta^b \partial_b N_a - \frac{1}{2} e_\alpha^a e_\beta^b N^c (\partial_b h_{ac} + \partial_a h_{bc} - \partial_c h_{ab}), \quad (7.28)$$

where the first term $e_\alpha^a e_\beta^b \partial_b N_a$ is identical on the two sides of F_L , and the remaining terms are of course zero on the Minkowski side F_L^+ . Because $h_{ab} = 0$ on F_L^- the derivatives $e_\alpha^a \partial_a h_{bc}$ tangent to F_L^- are all zero, but the limiting values of the transverse derivatives $N^c \partial_c h_{ab}$ are not necessarily zero there. So

$$[C_{\alpha\beta}] = -\frac{1}{2} e_\alpha^a e_\beta^b N^c \partial_c h_{ab}|_{F_L^-}. \quad (7.29)$$

Now, the derivatives of h_{ab} can be evaluated formally by using (5.22) to write:

$$\partial_c h_{ab}(t, \mathbf{x}) = [(\frac{d}{d\psi} R_a^p) h_{pq}(\bar{t}, \bar{\mathbf{x}}) R_b^q + R_a^p h_{pq}(\bar{t}, \bar{\mathbf{x}}) (\frac{d}{d\psi} R_b^q)] \partial_c \psi + (\partial \bar{x}^r / \partial x^c) R_a^p \bar{\partial}_r h_{pq}(\bar{t}, \bar{\mathbf{x}}) R_b^q, \quad (7.30)$$

where $\partial_c \psi$ and $\partial \bar{x}^r / \partial x^c$ can in turn be calculated from (5.12) and (5.13). However, as is shown in Appendix A.3, the derivatives $\bar{\partial}_r h_{pq}(\bar{t}, \bar{\mathbf{x}})$ are divergent at image points that lie in the beaming directions of the two kinks, and the loci of such points extend to future null infinity on F_0 . Although points of this type have $\bar{z} = \pm \frac{1}{8} L_0$, and so (since $z = e^{-\kappa \mu \psi} \bar{z}$) the only points on F_L which are limit points

of the singularities of $\partial_c h_{ab}$ lie on the plane $z = 0$, this feature of the metric makes (7.30) less than ideal for calculating $[C_{\alpha\beta}]$.

Instead, it is more convenient when calculating $[C_{\alpha\beta}]$ to use the formal definition of the derivative

$$\lim_{\|\delta y\| \rightarrow 0} \|\delta y\|^{-1} [f(y^a + \delta y^a) - f(y^a) - \delta y^c \partial_c f] = 0, \quad (7.31)$$

where f denotes any of the relevant combinations of the components of h_{ab} , and $\|\cdot\|$ is the Euclidean norm. This approach has the advantage of demonstrating that the relevant combinations of the metric derivatives do in fact exist at all points on the null boundary (except at p_L itself).

Consider then a fixed point $y^a \equiv [t_L + r, r\hat{\mathbf{r}}]^a$ on $F_L \setminus \{p_L\}$ (with $\hat{\mathbf{r}} \equiv (\hat{x}, \hat{y}, \hat{z})$ and $|\hat{\mathbf{r}}| = 1$), and let $x^a = y^a + \delta y^a$ be a field point at a Euclidean distance ϖt_L from (and in the timelike past of) y^a , so that $\delta y^a \equiv \varpi t_L [\delta \hat{t}, \delta \hat{\mathbf{r}}]^a$ where $\delta \hat{\mathbf{r}} \equiv (\delta \hat{x}, \delta \hat{y}, \delta \hat{z})$ and $\|[\delta \hat{t}, \delta \hat{\mathbf{r}}]\| = 1$. If ε denotes $e^{-\kappa\mu\psi}$ (as in the previous section) then

$$\varepsilon = |t_L^{-1} r \hat{\mathbf{r}} + \varpi \delta \hat{\mathbf{r}}| - t_L^{-1} r - \varpi \delta \hat{t} = (\hat{\mathbf{r}} \cdot \delta \hat{\mathbf{r}} - \delta \hat{t}) \varpi + O(\varpi^2). \quad (7.32)$$

It is easily seen that $\varepsilon \leq \sqrt{2}\varpi$, and that $\varepsilon > 0$ when x^a lies in the past of y^a .

The image point $\bar{x}^a$ corresponding to x^a has components

$$\begin{aligned} \bar{t} &= \varepsilon^{-1} |r \hat{\mathbf{r}} + \varpi t_L \delta \hat{\mathbf{r}}|, & \bar{x} &= \varepsilon^{-1} [(r \hat{x} + \varpi t_L \delta \hat{x}) \cos \psi + (r \hat{y} + \varpi t_L \delta \hat{y}) \sin \psi], \\ \bar{y} &= \varepsilon^{-1} [(r \hat{y} + \varpi t_L \delta \hat{y}) \cos \psi - (r \hat{x} + \varpi t_L \delta \hat{x}) \sin \psi] & \text{and} & \quad \bar{z} = \varepsilon^{-1} (r \hat{z} + \varpi t_L \delta \hat{z}). \end{aligned} \quad (7.33)$$

If the non-dimensionalized components $[T, X, Y, Z] = 4\pi L_0^{-1} [\bar{t}, \bar{x}, \bar{y}, \bar{z}]$ are substituted into the equation (5.33) for V_k then, after squaring both sides and rearranging, the equation reads (for $k = 0$ or 1)

$$|r \hat{\mathbf{r}} + \varpi t_L \delta \hat{\mathbf{r}}| (V_k + k\pi) - r \hat{\mathbf{r}} \cdot \mathbf{X}_k = \varpi t_L \left\{ \delta \hat{\mathbf{r}} \cdot \mathbf{X}_k + \frac{1}{2} \kappa \mu (\varepsilon / \varpi) [(V_k + k\pi)^2 - (1 + \pi^2/4)] \right\}, \quad (7.34)$$

where the factor of $\kappa\mu$ arises because $\frac{1}{4\pi} L_0 = \kappa\mu t_L$, and

$$\mathbf{X}_k \equiv (\cos(V_k + \psi), \sin(V_k + \psi), (k - \frac{1}{2})\pi). \quad (7.35)$$

Equation (7.34) suggests that, for small values of ϖ , the root V_k can be approximated by the solution of the equation $V_k + k\pi = \hat{\mathbf{r}} \cdot \mathbf{X}_k$, which I will denote by V_k^∞ . In explicit form, the equations for $V_{0,1}^\infty$ read

$$V_0^\infty = \hat{x} \cos(V_0^\infty + \psi) + \hat{y} \sin(V_0^\infty + \psi) - \frac{\pi}{2} \hat{z} \quad (7.36)$$

and

$$V_1^\infty + \pi = \hat{x} \cos(V_1^\infty + \psi) + \hat{y} \sin(V_1^\infty + \psi) + \frac{\pi}{2} \hat{z}. \quad (7.37)$$

Note however that $V_{0,1}^\infty$ are typically still functions of ϖ , as $\psi = -(\kappa\mu)^{-1} \ln \varepsilon$ increases without bound, and in particular $\lim_{\varpi \rightarrow 0} V_k^\infty$ does not exist unless $\hat{x} = \hat{y} = 0$.

Although $\lim_{\varpi \rightarrow 0} V_k^\infty$ does not generally exist, it can be shown that $\lim_{\varpi \rightarrow 0} |V_k - V_k^\infty| = 0$, a fact that will prove important in establishing that certain combinations of the metric components are differentiable on F_L . To show that $\lim_{\varpi \rightarrow 0} |V_k - V_k^\infty| = 0$, first rearrange (7.34) to give

$$\begin{aligned} r(V_k + k\pi - \hat{\mathbf{r}} \cdot \mathbf{X}_k) &= \varpi t_L \{ \delta \hat{\mathbf{r}} \cdot \mathbf{X}_k + \frac{1}{2} \kappa \mu (\varepsilon / \varpi) [(V_k + k\pi)^2 - (1 + \pi^2/4)] \} \\ &\quad + (r - |r\hat{\mathbf{r}} + \varpi t_L \delta \hat{\mathbf{r}}|)(V_k + k\pi) \\ &\equiv \Theta(\varpi). \end{aligned} \quad (7.38)$$

Since $|V_k + k\pi|$ is bounded above (see Appendix A.1), it is clear that the function $\Theta(\varpi)$ on the right-hand side of this equation tends to zero as $\varpi \rightarrow 0$. The root V_k^∞ satisfies the same equation (7.38) with $\Theta(\varpi) = 0$.

Subtracting the equation for V_k^∞ from (7.38) gives:

$$V_k - V_k^\infty - \hat{x} [\cos(V_k + \psi) - \cos(V_k^\infty + \psi)] - \hat{y} [\sin(V_k + \psi) - \sin(V_k^\infty + \psi)] = r^{-1} \Theta(\varpi). \quad (7.39)$$

In view of the identities $\cos 2A - \cos 2B = 2 \sin(B - A) \sin(A + B)$ and $\sin 2A - \sin 2B = 2 \sin(A - B) \cos(A + B)$ this equation becomes

$$V_k - V_k^\infty + 2[\hat{x} \sin(\frac{1}{2}V_k + \frac{1}{2}V_k^\infty + \psi) - \hat{y} \cos(\frac{1}{2}V_k + \frac{1}{2}V_k^\infty + \psi)] \sin[\frac{1}{2}(V_k - V_k^\infty)] = r^{-1} \Theta(\varpi), \quad (7.40)$$

and so

$$|V_k - V_k^\infty| - 2(\hat{x}^2 + \hat{y}^2)^{1/2} |\sin[\frac{1}{2}(V_k - V_k^\infty)]| \leq r^{-1} |\Theta(\varpi)|. \quad (7.41)$$

Since $(\hat{x}^2 + \hat{y}^2)^{1/2} \leq 1$ and $|w| - 2|\sin(\frac{1}{2}w)| \geq \frac{1}{48}|w|^3$ for all real w , it follows that

$$\frac{1}{48} |V_k - V_k^\infty|^3 \leq r^{-1} |\Theta(\varpi)| \quad (7.42)$$

and therefore $\lim_{\varpi \rightarrow 0} |V_k - V_k^\infty| = 0$ as claimed. Furthermore, because $\Theta(\varpi)$ goes to zero linearly in ϖ , $|V_k - V_k^\infty|$ is at worst of order $\varpi^{1/3}$ for small values of ϖ .

The asymptotic forms of the components of h_{ab} can now be developed by first substituting the expressions

$$\chi_\pm = 4\pi L_0^{-1} \varepsilon^{-1} (t \pm z) \pm \frac{\pi}{2} \quad (7.43)$$

into (5.38)-(5.40) to give

$$\begin{aligned} I_+ &= \frac{1}{4\pi} L_0 \varepsilon \frac{V_0 - V_1 - 2\pi}{t+z} + O(\varepsilon^2), & I_- &= \frac{1}{4\pi} L_0 \varepsilon \frac{V_1 - V_0}{t-z} + O(\varepsilon^2), \\ S_\pm &= \pm \frac{1}{4\pi} L_0 \varepsilon \frac{\cos V_1 - \cos V_0}{t \pm z} + O(\varepsilon^2) & \text{and} & \quad C_\pm = \pm \frac{1}{4\pi} L_0 \varepsilon \frac{\sin V_0 - \sin V_1}{t \pm z} + O(\varepsilon^2) \end{aligned} \quad (7.44)$$

(assuming for the moment that $t \pm z \neq 0$).

With $t = |r\hat{\mathbf{r}} + \varpi t_L \delta\hat{\mathbf{r}}|$ and $z = r\hat{z} + \varpi t_L \delta\hat{z}$, equation (5.37) generates the following expansion for the components of h_{ab} :

$$\begin{aligned} h_{ab}(t, \mathbf{x}) &= R_a^c(\psi) h_{cd}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}}) R_b^d(\psi) \\ &= \frac{2L_0\mu\varepsilon}{\pi r(1-\hat{z}^2)} \begin{bmatrix} \Delta V \hat{z} - (1-\hat{z})\pi & (c_0 - c_1)\hat{z} & (s_0 - s_1)\hat{z} & -\Delta V - (1-\hat{z})\pi \\ (c_0 - c_1)\hat{z} & \Delta V \hat{z} - (1-\hat{z})\pi & 0 & c_1 - c_0 \\ (s_0 - s_1)\hat{z} & 0 & \Delta V \hat{z} - (1-\hat{z})\pi & s_1 - s_0 \\ -\Delta V - (1-\hat{z})\pi & c_1 - c_0 & s_1 - s_0 & \Delta V \hat{z} - (1-\hat{z})\pi \end{bmatrix}_{ab} \\ &\quad + O(\varpi^2), \end{aligned} \quad (7.45)$$

where

$$c_k \equiv \cos(V_k + \psi), \quad s_k \equiv \sin(V_k + \psi) \quad \text{and} \quad \Delta V \equiv V_1 - V_0. \quad (7.46)$$

In terms of the components of $\hat{\mathbf{r}}$ the relevant orthonull basis vectors are

$$k^a = [1, \hat{x}, \hat{y}, \hat{z}]^a, \quad e_1^a = r(1-\hat{z}^2)^{-1/2} [0, \hat{x}\hat{z}, \hat{y}\hat{z}, -(\hat{x}^2 + \hat{y}^2)] \quad \text{and} \quad e_2^a = r[0, -\hat{y}, \hat{x}, 0]^a, \quad (7.47)$$

and so

$$k^a k^b h_{ab} = -4(L_0/r)\mu\varepsilon + O(\varpi^2), \quad k^a e_2^b h_{ab} = O(\varpi^2) \quad (7.48)$$

$$k^a e_1^b h_{ab} = \frac{2}{\pi(1-\hat{z}^2)^{3/2}} L_0 \mu \varepsilon (V_1 + \pi - \hat{\mathbf{r}} \cdot \mathbf{X}_1 - V_0 + \hat{\mathbf{r}} \cdot \mathbf{X}_0) + O(\varpi^2), \quad (7.49)$$

and

$$\begin{aligned} \sigma^{AB} e_A^a e_B^b h_{ab} &= -r^{-2} [e_1^a e_1^b h_{ab} + (1-\hat{z}^2)^{-1} e_2^a e_2^b h_{ab}] \\ &= -\frac{4}{\pi(1-\hat{z}^2)} (L_0/r) \mu \varepsilon [(V_1 + \pi - \hat{\mathbf{r}} \cdot \mathbf{X}_1 - V_0 + \hat{\mathbf{r}} \cdot \mathbf{X}_0) \hat{z} - \pi(1-\hat{z}^2)] \\ &\quad + O(\varpi^2). \end{aligned} \quad (7.50)$$

Now, given that $\lim_{\varpi \rightarrow 0} |V_k - V_k^\infty| = 0$ and c_k and s_k are continuous functions of V_k ,

$$\lim_{\varpi \rightarrow 0} (V_1 + \pi - \hat{\mathbf{r}} \cdot \mathbf{X}_1) = \lim_{\varpi \rightarrow 0} [V_1^\infty + \pi - \hat{x} \cos(V_1^\infty + \psi) - \hat{y} \sin(V_1^\infty + \psi) - \frac{\pi}{2} \hat{z}] = 0, \quad (7.51)$$

and similarly $\lim_{\varpi \rightarrow 0}(V_0 - \hat{\mathbf{r}} \cdot \mathbf{X}_0) = 0$. Hence, in view of the expansion (7.32),

$$\begin{aligned} \lim_{\varpi \rightarrow 0} \varpi^{-1} k^a k^b h_{ab} &= -4(L_0/r) \mu (\hat{\mathbf{r}} \cdot \delta \hat{\mathbf{r}} - \delta \hat{t}), & \lim_{\varpi \rightarrow 0} \varpi^{-1} k^a e_1^b h_{ab} &= 0 \\ \lim_{\varpi \rightarrow 0} \varpi^{-1} k^a e_2^b h_{ab} &= 0 & \text{and} & \lim_{\varpi \rightarrow 0} \varpi^{-1} \sigma^{AB} e_A^a e_B^b h_{ab} &= 4(L_0/r) \mu (\hat{\mathbf{r}} \cdot \delta \hat{\mathbf{r}} - \delta \hat{t}). \end{aligned} \quad (7.52)$$

Since $\delta y^a \equiv \varpi t_L [\delta \hat{t}, \delta \hat{\mathbf{r}}]^a$ it follows from (7.31) that $k^a k^b h_{ab}$, $k^a e_A^b h_{ab}$ and $\sigma^{AB} e_A^a e_B^b h_{ab}$ are all differentiable at each point on the null boundary F_L^- with $\hat{z}^2 \neq 1$, and that

$$k^a k^b \partial_c h_{ab} = -\sigma^{AB} e_A^a e_B^b \partial_c h_{ab} = 4(L_0/r) t_L^{-1} \mu [1, -\hat{\mathbf{r}}]_c \quad \text{and} \quad k^a e_A^b \partial_c h_{ab} = 0 \quad (7.53)$$

there. Given that $N^c[1, -\hat{\mathbf{r}}]_c = 1$, it seems therefore that

$$[C_{33}] = -\sigma^{AB} [C_{AB}] = -2(L_0/r) t_L^{-1} \mu \quad \text{and} \quad \sigma^{AB} [C_{3B}] = 0, \quad (7.54)$$

and hence that matching the evaporating ACO spacetime to the Minkowski spacetime across F_L induces a boundary layer with a non-zero density ρ and isotropic pressure p .

However, it should be recalled that $L_0 t_L^{-1} = 4\pi \kappa \mu$, and so the apparent density and pressure of this boundary layer,

$$\rho = -p = -\kappa \mu^2 / r, \quad (7.55)$$

are quadratic in μ .⁸ Thus ρ and p are zero at the level of the first-order approximation used throughout this paper, and the evaporating ACO spacetime can be matched to the Minkowski spacetime without inducing a boundary layer *at this level of approximation*.

All that remains now is to show that the same conclusion holds at all points on F_L^- with $\hat{z}^2 = 1$ (that is, on the z -axis). In this case, $\hat{x} = \hat{y} = 0$ and V_k tends to the limiting value

$$V_k^\infty = -k\pi + (k - \frac{1}{2})\pi \hat{z}. \quad (7.56)$$

Also, it is convenient to use a slightly different asymptotic form for the spatial components of the field point x^a , namely

$$x = \frac{1}{4\pi} L_0 x^* \varepsilon, \quad y = \frac{1}{4\pi} L_0 y^* \varepsilon \quad \text{and} \quad z = r \hat{z} + \frac{1}{4\pi} L_0 z^* \varepsilon, \quad (7.57)$$

⁸Incidentally, it is to be expected that neither ρ nor p depend separately on L_0 or t_L , as the values of the latter will vary as the zero point of t changes, but the stress-energy content of F_L should remain invariant under time translations.

where x^* , y^* and z^* are all of order ε^0 .

Then

$$X = x^* \cos \psi + y^* \sin \psi, \quad Y = y^* \cos \psi - x^* \sin \psi \quad \text{and} \quad Z = 4\pi L_0^{-1} \varepsilon^{-1} r \hat{z} + z^*, \quad (7.58)$$

and the analogue of (7.43) becomes

$$\chi_{\pm} = 4\pi r L_0^{-1} (1 \pm \hat{z}) \varepsilon^{-1} + z^* \hat{z} (1 \pm \hat{z}) \pm \frac{1}{2} \pi + \frac{1}{8\pi} L_0 r^{-1} (X^2 + Y^2) \varepsilon - \frac{1}{2} \left(\frac{1}{4\pi} L_0 \right)^2 r^{-2} z^* \hat{z} (X^2 + Y^2) \varepsilon^2 \quad (7.59)$$

to order ε^2 , while the analogue of equation (7.38) for V_k can be used to calculate the asymptotic expansion

$$\begin{aligned} \chi_{\pm} - V_k &= 4\pi r L_0^{-1} (1 \pm \hat{z}) \varepsilon^{-1} + z^* \hat{z} (1 \pm \hat{z}) \pm \frac{1}{2} \pi + k\pi - (k - \frac{1}{2}) \pi \hat{z} \\ &\quad + \frac{1}{8\pi} L_0 r^{-1} (X^2 + Y^2 + 2Y \hat{z} + 1) \varepsilon + \frac{1}{2} \left(\frac{1}{4\pi} L_0 \right)^2 r^{-2} X (2Y + \hat{z}) \varepsilon^2 \\ &\quad + \frac{1}{2} \left(\frac{1}{4\pi} L_0 \right)^2 r^{-2} \hat{z} [(k - \frac{1}{2}) \pi - z^*] (X^2 + Y^2 + 2Y \hat{z} + 1) \varepsilon^2 \end{aligned} \quad (7.60)$$

(plus terms of order ε^3).

The expansions need to be developed to such a high order because if $\hat{z} = 1$ (*resp.* $\hat{z} = -1$) then the terms appearing at order ε^{-1} and ε^0 make no contribution to I_- , S_- or C_- (*resp.* I_+ , S_+ or C_+), and the terms of order ε plus the terms proportional to $(k - \frac{1}{2})$ at order ε^2 dominate the asymptotic behavior.. Otherwise (if $\hat{z} = -1$, *resp.* $\hat{z} = 1$), it is the terms at order ε^{-1} and ε^0 that dominate. A straightforward but lengthy calculation shows that

$$\begin{aligned} I_+ &= I_- = -\frac{1}{4} (L_0/r) \varepsilon, & S_+ &= -\frac{1}{8} (1 - \hat{z}) (L_0/r) \varepsilon, \\ S_- &= \frac{1}{8} (1 + \hat{z}) (L_0/r) \varepsilon & \text{and} & \quad C_+ = C_- = 0 \end{aligned} \quad (7.61)$$

to order ε , and hence from (5.37) that

$$h_{ab}(t, \mathbf{x}) = (L_0/r) \mu \varepsilon \begin{bmatrix} -2 & \hat{z} \cos \psi & \hat{z} \sin \psi & 0 \\ \hat{z} \cos \psi & -2 & 0 & -\cos \psi \\ \hat{z} \sin \psi & 0 & -2 & -\sin \psi \\ 0 & -\cos \psi & -\sin \psi & -2 \end{bmatrix}_{ab} + O(\varepsilon^2). \quad (7.62)$$

Furthermore, when $\hat{z}^2 = 1$ the spherical coordinates (r, θ^1, θ^2) parametrizing the null surface F_L are singular, as σ_{AB} is degenerate. A more suitable parametrization of F_L in this case would align the poles of the spherical coordinate system with (say) the x -axis. Then

$$k^a = [1, 0, 0, \hat{z}]^a, \quad e_1^a = r[0, 0, -1, 0] \quad \text{and} \quad e_2^a = r[0, \hat{z}, 0, 0]^a, \quad (7.63)$$

and $\sigma^{AB} = -r^{-2}\delta^{AB}$. With this choice of orthonull basis, it is easily checked that

$$k^a k^b h_{ab} = -4(L_0/r)\mu\varepsilon + O(\varepsilon^2), \quad k^a e_1^b h_{ab} = O(\varepsilon^2), \quad k^a e_2^b h_{ab} = O(\varepsilon^2) \quad (7.64)$$

and

$$\sigma^{AB} e_A^a e_B^b h_{ab} = 4(L_0/r)\mu\varepsilon + O(\varepsilon^2), \quad (7.65)$$

just as in the case $\hat{z}^2 \neq 1$ considered earlier. So it can be seen that the null boundary $F_L \setminus \{p_L\}$ separating the evaporating ACO spacetime from the relic Minkowski spacetime contains no stress-energy to first order in μ .

8. Dynamical Stability of the ACO Loop

Realistic cosmic string loops would not, of course, have the exact shape of an ACO loop, and would not evaporate in a strictly self-similar manner. It is therefore important to be able to estimate the impact that deviations from the ACO loop trajectory would have on the evaporation process. A full analysis of the stability of the ACO loop in the presence of back-reaction is beyond the scope of this paper (not least because the dynamical consequences of the self-gravity of a string loop in the strong-field limit are not yet understood), and in this section I will limit myself to some observations about the dynamical stability of the flat-space ACO solution.

The trajectory of the flat-space ACO loop is fully described by the position vector $\mathbf{X}(\tau, \sigma)$ defined in (3.2), where the mode functions $\mathbf{a}$ and $\mathbf{b}$ are specified by equations (3.11) and (3.12). For the purposes of the perturbation analysis of this section I will refer to the unperturbed mode functions as $\mathbf{a}_0$ and $\mathbf{b}_0$, so that

$$\mathbf{a}_0(\sigma_+) = (|\sigma_+| - \tfrac{1}{4}L) \hat{\mathbf{z}} \quad \text{for} \quad -\tfrac{1}{2}L \leq \sigma_+ \leq \tfrac{1}{2}L \quad (8.1)$$

and

$$\mathbf{b}_0(\sigma_-) = \frac{L}{2\pi} [\cos(2\pi\sigma_-/L) \hat{\mathbf{x}} + \sin(2\pi\sigma_-/L) \hat{\mathbf{y}}], \quad (8.2)$$

where $\sigma_{\pm} = \tau \pm \sigma$. (Strictly speaking, of course, $\mathbf{a}_0$ is the even periodic extension of the function on the right of (8.1).)

A generic perturbation of the ACO solution has mode functions of the form

$$\mathbf{a}(\sigma_+) = \mathbf{a}_0(\sigma_+) + \delta\mathbf{a}(\sigma_+) \quad \text{and} \quad \mathbf{b}(\sigma_-) = \mathbf{b}_0(\sigma_-) + \delta\mathbf{b}(\sigma_-), \quad (8.3)$$

where $\delta\mathbf{a}$ and $\delta\mathbf{b}$ are continuous functions of their arguments, and are both small, in the sense that $|\delta\mathbf{a}| \ll L$ and $|\delta\mathbf{b}| \ll L$ everywhere. Without loss of generality it

can be assumed that the perturbed solution has the same parametric period, L , as the unperturbed ACO loop, and so $\delta\mathbf{a}$ and $\delta\mathbf{b}$ are periodic functions with period L . The only other constraints on $\delta\mathbf{a}$ and $\delta\mathbf{b}$ are the gauge conditions $|\mathbf{a}'|^2 = |\mathbf{b}'|^2 = 1$, which to linear order in the perturbations require that $\mathbf{a}'_0 \cdot \delta\mathbf{a}' = 0$ and $\mathbf{b}'_0 \cdot \delta\mathbf{b}' = 0$ at all points where these derivatives exist.

Note in particular that the derivatives $\delta\mathbf{a}'$ and $\delta\mathbf{b}'$ need not be continuous, nor is it necessarily true that either $|\delta\mathbf{a}'|$ or $|\delta\mathbf{b}'|$ is small. A simple way to visualize the effect of a perturbation is to plot the unit vectors $\mathbf{a}'$ and $\mathbf{b}'$ on the surface of a unit sphere, an approach first popularized by Kibble and Turok [22]. Figure 6 contains three plots of this type. The first represents the unperturbed ACO loop, and shows the $\mathbf{b}'$ curve following the equator of the sphere while $\mathbf{a}' = \pm\hat{\mathbf{z}}$ is concentrated at just two points, the north and south poles. In the second plot, the perturbations $\delta\mathbf{a}'$ and $\delta\mathbf{b}'$ are assumed to be both continuous and small, with the result that the $\mathbf{a}'$ and $\mathbf{b}'$ curves depart only minimally from the unperturbed curves. The $\mathbf{a}'$ curve, in particular, is understood to jump discontinuously from the small circle around the south pole to the small circle around the north pole when $\sigma_+ \approx 0$ then back again when $\sigma_+ \approx \pi$, so the two kinks are preserved under perturbations of this form.

The third plot in Figure 6 shows the effect of a perturbation in which $\delta\mathbf{a}'$ is continuous but no longer small. In fact, $|\delta\mathbf{a}'|$ is assumed to be of order unity for a narrow range of values of the parameter σ_+ centered on $\sigma_+ \approx 0$ and $\sigma_+ \approx \pi$, creating two smooth segments on the $\mathbf{a}'$ curve linking the north and south poles. This has the effect of removing the discontinuity in the $\mathbf{a}'$ curve, and thus smoothing the two kinks, but an unavoidable consequence is that the $\mathbf{a}'$ and $\mathbf{b}'$ curves now cross at two points, indicating that a pair of cusps will momentarily form during each oscillation of the perturbed loop.

So although the unperturbed ACO loop does not support cusps, it is evident that there exist solutions of the flat-space equation of motion arbitrarily close to the ACO loop in $\mathbf{a}$ - $\mathbf{b}$ trajectory space that do support cusps. The presence of cusps complicates the back-reaction problem considerably. Because the bulk velocity of a string loop momentarily reaches the speed of light c at a cusp, a non-negligible fraction of the loop's total energy can be concentrated there. A simple kinematic argument ([23], Section 6.2) suggests that the total energy M_r inside a sphere of radius r centered on a cusp scales as $r^{1/2}$, and so the gravitational potential M_r/r diverges as $r^{-1/2}$. This in turn is an indication that the effects of gravitational back-reaction at a cusp cannot be modeled accurately at the level

of the weak-field approximation.⁹

Any cusps forming on a perturbed solution that is close to the ACO loop will necessarily be very narrow, as the energy in such a cusp will be proportional to the range of the parameter σ_+ covered by the segment linking the two poles on the Kibble-Turok sphere. What effect back-reaction would have on the cusps is still an open question. It is possible that back-reaction would act to narrow the cusps even further, thus driving the perturbed solution closer to the unperturbed ACO loop. The ACO loop would then be dynamically stable.

Alternatively, the cusps could broaden in the presence of back-reaction and ultimately expand to the size of ordinary macrocusps, which typically contain a fraction of order μ of the total energy of the string loop ([23], Section 6.2). In this case, it seems likely that the perturbed loop will evaporate very quickly, as cusps are known to beam away copious amounts of gravitational radiation [19]. Indeed, the presumption alluded to in Section 3 that the ACO solution has the lowest radiative efficiency of any cosmic string loop encourages the view that dynamical instabilities in the trajectory of the ACO loop, if they exist, are likely to be of only secondary importance. A perturbed trajectory that diverges rapidly from the ACO loop will have an increased radiative efficiency and will evaporate more quickly, whereas perturbations that remain close to the ACO loop will survive longer.

A third possibility is that cusps on perturbed solutions close to the ACO loop are unstable to gravitational collapse, with consequences that have not yet been explored and can at present only be imagined.

Another potential source of dynamical instability is loop self-intersection. String loops that intersect themselves are believed to split into two daughter loops with high probability ([23], Section 2.7). The ACO loop itself has no self-intersections, but there do exist trajectories arbitrarily close to the ACO loop that intersect themselves, as will be seen shortly.

A self-intersection occurs on a general flat-space loop when

$$\mathbf{a}(\sigma_+ + \Delta) - \mathbf{a}(\sigma_+) = \mathbf{b}(\sigma_-) - \mathbf{b}(\sigma_- - \Delta) \quad (8.4)$$

for some values of σ_+ , σ_- and Δ , with $\Delta \in (0, L/2)$ ([23], Section 3.8). If $\mathbf{a}$ and $\mathbf{b}$ are decomposed in accordance with (8.3), the x - and y -components of equation

⁹By contrast, because the bulk Lorentz factor of a string is bounded at a kink, the energy M_r inside a sphere of radius r centered on a kink typically scales as r , and the gravitational potential M_r/r has no problematic divergences. The breakdown of the weak-field approximation at a kink is due not to a local spike in the energy content of the string, but to the fact that the tangent vector $X_{(0),u}^a$ is discontinuous (see Section 9 below).

(8.4) read:

$$\cos(2\pi\sigma_-/L) - \cos[2\pi(\sigma_- - \Delta)/L] = 2\pi L^{-1}(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot \hat{\mathbf{x}} \quad (8.5)$$

and

$$\sin(2\pi\sigma_-/L) - \sin[2\pi(\sigma_- - \Delta)/L] = 2\pi L^{-1}(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot \hat{\mathbf{y}} \quad (8.6)$$

respectively, where

$$\delta_\Delta \mathbf{a} \equiv \delta \mathbf{a}(\sigma_+ + \Delta) - \delta \mathbf{a}(\sigma_+) \quad \text{and} \quad \delta_\Delta \mathbf{b} \equiv \delta \mathbf{b}(\sigma_-) - \delta \mathbf{b}(\sigma_- - \Delta). \quad (8.7)$$

Equations (8.5) and (8.6) can be solved to give:

$$\cos(2\pi\sigma_-/L) = \pi L^{-1}(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot [\hat{\mathbf{x}} + \frac{\sin(2\pi\Delta/L)}{1 - \cos(2\pi\Delta/L)} \hat{\mathbf{y}}] \quad (8.8)$$

and

$$\sin(2\pi\sigma_-/L) = \pi L^{-1}(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot [\hat{\mathbf{y}} - \frac{\sin(2\pi\Delta/L)}{1 - \cos(2\pi\Delta/L)} \hat{\mathbf{x}}]. \quad (8.9)$$

In particular,

$$1 - \cos(2\pi\Delta/L) = 2\pi^2 L^{-2} [|(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot \hat{\mathbf{x}}|^2 + |(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot \hat{\mathbf{y}}|^2], \quad (8.10)$$

and so Δ must be of order $|\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}|$ or smaller. For small perturbations of the ACO loop, therefore, self-intersections can only occur with $\Delta \ll L$.

Moreover, for small values of Δ equations (8.5) and (8.6) reduce to

$$\Delta^{-1}(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot \hat{\mathbf{x}} \approx -\sin(2\pi\sigma_-/L) \quad \text{and} \quad \Delta^{-1}(\delta_\Delta \mathbf{a} - \delta_\Delta \mathbf{b}) \cdot \hat{\mathbf{y}} \approx \cos(2\pi\sigma_-/L) \quad (8.11)$$

respectively, and can be satisfied only if either $\delta \mathbf{a}'$ or $\delta \mathbf{b}'$ is discontinuous or of order unity at the self-intersection point $(\sigma_+, \sigma_-) \approx (\sigma_+ + \Delta, \sigma_- - \Delta)$. Hence, at least one of the $\mathbf{a}'$ and $\mathbf{b}'$ curves will deviate significantly from the corresponding unperturbed ACO curve on the Kibble-Turok sphere.

The picture that emerges, therefore, is that a flat-space solution close to the ACO loop can intersect itself only by pinching off a small daughter loop with parametric period $\Delta \ll L$. A solution that does just this is easily constructed by choosing (at time $t = 0$ say) a small segment of the unperturbed ACO loop with length of order Δ and deforming it so that it crosses itself. But the same perturbation procedure could be applied to any flat-space loop solution, so self-intersections of this type are in a sense rather trivial.

Furthermore, the larger of the two daughter loops created by the self-intersection will be little changed from the original parent loop. It will have a marginally

smaller parametric length $L - \Delta$, and in place of the self-intersection there will be two narrow kinks traveling in opposite directions around the loop. Analytic studies of kinks on long strings strongly suggest that gravitational back-reaction will suppress small-scale structure on a string with a characteristic length r on a timescale $\Delta t \sim r/(8\pi^2\mu)$ ([23], Section 6.7), and this estimate has been confirmed in numerical simulations of planar loops carrying up to 32 kinks by Quashnock and Spergel [11]. In the case of a kink formed by the pinching off of a daughter loop with a parametric period Δ the characteristic length r will be of order Δ , and the decay time Δt for the kink will be much smaller than the radiative lifetime $t_L = L/(4\pi\kappa\mu)$ of the ACO loop. So it is to be expected that the two narrow kinks created by the self-intersection will quickly dissipate, leaving behind a daughter loop that very closely resembles the ACO loop.

To summarize, there is as yet no reason to believe that the ACO solution is dynamically unstable, although a definitive statement on the matter will depend on the results of full back-reaction stability analyses, both at the weak-field level (without cusps) and in the strong-field limit (with cusps). Nonetheless, it seems safe to conclude that the effect of possible self-intersections on the stability of the ACO loop is negligible. Furthermore, even if it were to turn out that the ACO loop is unstable to certain mechanisms (such as cusp formation), this would not in itself invalidate the importance of the self-similar evaporating ACO solution described above, as the longest-lived loops in any ensemble of loops with the same initial energy would be those that most closely resemble the ACO loop – on the presumption, of course, that the ACO solution is unique in having the lowest radiative efficiency of any string loop.

I should add that at present there is an ongoing debate about the nature of the scaling peaks in the production spectrum of cosmic string loops, fueled by the conflicting results of recent numerical [24, 25] and analytic [26, 27] studies of the primordial string network. However, this debate has no direct bearing on the stability or evolution of the ACO loop.

9. Conclusions

In this paper I have constructed a solution of the linearized Einstein and Nambu-Goto equations describing a cosmic string loop that evaporates self-similarly under the action of its own self-gravity, eventually radiating away all its energy and momentum to leave a remnant flat spacetime, and whose spacelike cross-sections are identical (to leading order in the string’s mass per unit length μ) to those of

the Allen-Casper-Ottewill loop.

From a purely physical viewpoint, the ACO loop is one of the most important flat-space cosmic string solutions, as it is long-lived and approximates well the lowest-efficiency daughter loops observed in low-resolution string network simulations [18, 19]¹⁰. The fact that its evolution and ultimate evaporation are analytically tractable is an added bonus, and makes the ACO loop an object of potentially great interest to mathematical relativity.

However, much work needs to be done before it could be claimed with any confidence that the evaporating ACO loop can be described by a self-consistent solution of the Einstein equations. A first step in this direction would be to construct a solution to the *strong-field* back-reaction problem that reduces to the bare ACO loop in the limit $\mu \rightarrow 0$ – or, at the very least, to demonstrate that such a solution exists. A second desideratum would be to find a solution of the Abelian Higgs field equations that can act as a suitable source for the evaporating ACO metric. Neither task is likely to be easy.

One question that arises naturally in this context is whether the self-similarity of the linearized solution would be preserved in the corresponding fully non-linear solution (if indeed such a solution exists). This question remains an open one, but it can be given a more rigorous formulation as follows. If the original Minkowski coordinates $x^a = [t, \mathbf{x}]^a$ describing the linearized solution are written in terms of the similarity coordinates $[\psi, \bar{\mathbf{x}}]$, where ψ and $\bar{\mathbf{x}}$ are defined by equations (5.12) and (5.13) respectively, then it can be shown that the vector field

$$k^a \equiv \partial x^a / \partial \psi = [0, -y, x, 0]^a - \kappa \mu [t - t_L, x, y, z]^a \quad (9.1)$$

satisfies the equation

$$(\eta_{c(a} + h_{c(a})\partial_b)k^c + \frac{1}{2}k^c\partial_c h_{ab} = -\kappa\mu\eta_{ab} \quad (9.2)$$

¹⁰More recent high-resolution simulations indicate that string loops, when they first form in an evolved string network, have an approximately fractal structure, with a fractal dimension approaching 2 on large scales [28, 29, 30]. The ACO loop therefore approximates realistic incipient string loops less closely than was first believed. Nonetheless, it remains true that the ACO loop has the lowest known radiative efficiency of any string loop. Furthermore, as was explained in Section 8, there are good theoretical reasons for believing that radiative back-reaction acts to preferentially eliminate small-scale structure from a loop on timescales much shorter than the lifetime of the loop (although the principal decay channel for structure on the very smallest scales would probably be particle production [28]). Hence, it seems likely that any fractal loop similar in shape to an ACO loop on large scales would radiate away much of its small-scale structure and quickly come to resemble the ACO loop on all scales. It has also been suggested [26, 27] that loop fragmentation will quickly act to suppress fractal structure, by triggering a cascade of daughter loops with progressively fewer cusps and kinks.

to linear order in μ . That is, k^a satisfies the linearized version of the equation $\nabla_{(a}k_{b)} = -\kappa\mu g_{ab}$ for a conformal Killing vector field.

The corresponding non-linear solution will therefore preserve self-similarity if it admits a vector field k^a satisfying the *exact* conformal Killing equation $\nabla_{(a}k_{b)} = -\kappa\mu g_{ab}$, in which case it is always possible to find coordinates $[\psi, \bar{\mathbf{x}}]$ in terms of which the metric tensor has the conformal form

$$g_{ab}(\psi, \bar{\mathbf{x}}) = e^{-2\kappa\mu\psi} p_{ab}(\bar{\mathbf{x}}). \quad (9.3)$$

(The conformal Killing field in this case is $k^a = \delta^a_\psi$.) So the question of whether self-similarity would be preserved by a fully non-linear solution can be recast as a question about the existence of solutions to the strong-field back-reaction problem with the conformal form (9.3).

Another of the more problematic aspects of the ACO loop is the behavior of the gravitational field near the two kink points. The discontinuity in the null tangent vector $X^a_{(0),u}$ at the kinks generates two apparent anomalies in the weak-field analysis described in Section 6. The first is that the term $X^d_{(0),CD}$ appearing in the first-order equation of motion (6.28) is undefined at the kinks, and so it cannot meaningfully be said that the equation of motion is satisfied at these points. The second anomaly is that the metric perturbations h_{ab} fail to be differentiable on a manifold of points extending from the trajectories of the kinks out to future null infinity, as is explained in more detail in Appendix A.3.

The breakdown in the differentiability of h_{ab} poses no intractable difficulty for the linearized analysis of Section 6, because the ill-defined terms make no overall contribution to the *linearized* Einstein tensor. However, the fully non-linear Einstein tensor contains products of terms of the form $\partial_c g_{ab}$, and in the generic case it is expected that a distributional interpretation of the Einstein equation is possible only if the metric derivatives $\partial_c g_{ab}$ are locally square integrable [2]. So the breakdown of differentiability in the beaming directions of the kinks may bode ill for the prospects of a strong-field solution.

A further problem is that the non-differentiability of h_{ab} is superimposed on the conical singularity marking the string worldsheet at the kink points themselves. The singularity in h_{ab} at the kinks was examined briefly in Paper I, and is far more pathological than the simple logarithmic singularity that appears at all other points on the string worldsheet. An appreciation for the nature of the singularity can be gained by repeating the analysis performed earlier, in Section 6.2, to calculate the near-field metric perturbations at a kink point.

If the field point $x^a = \frac{1}{4\pi}L_0[T, X, Y, Z]^a$ is chosen to be close to the lower of the kink points (at $u = 0$) near the spacelike surface $t = 0$, so that

$$[T, X, Y, Z]^a = [v, \cos v, \sin v, -\frac{1}{2}\pi]^a + 4\pi L_0^{-1}\delta x^a \quad (9.4)$$

for some $v \in (-\pi, \pi)$, then it can be shown that $\chi_{\pm} = v + 4\pi L_0^{-1}(\delta t \pm \delta z)$, and that

$$\begin{aligned} \chi_+ - V_1 - 2\pi &= \frac{1}{2\pi}(4\pi L_0^{-1})^2 Q_-, & \chi_- - V_1 &= 2\pi - 8\pi L_0^{-1}\delta z \\ \text{and} \quad \chi_{\pm} - V_0 &= 4\pi L_0^{-1}(\delta t \pm \delta z - \frac{1}{2} \frac{\eta_{ab}\delta x^a\delta x^b}{\delta t + \delta x \sin v - \delta y \cos v}) \end{aligned} \quad (9.5)$$

to first or leading order in δx^a , where

$$\begin{aligned} Q_- &\equiv (\delta x)^2 + (\delta y)^2 + (\delta t + \delta z)(\delta t + \delta z + 2\delta x \sin v - 2\delta y \cos v) \\ &\equiv \bar{\Psi}_{ab}(-1, v) \delta x^a \delta x^b. \end{aligned} \quad (9.6)$$

All four of the functions in (9.5) appear as arguments of the logarithmic and sine and cosine integral functions contributing to the terms $I_{\pm}$, $S_{\pm}$ and $C_{\pm}$ in (5.37). For points on the future lightcone of the kink, where $\eta_{ab}\delta x^a\delta x^b = 0$, the expansions (9.5) simplify considerably, as $\chi_{\pm} - V_0$ then becomes $4\pi L_0^{-1}(\delta t \pm \delta z)$, while $Q_- = 2(\delta t + \delta z)(\delta t + \delta x \sin v - \delta y \cos v)$. (Furthermore, the function $\delta t + \delta x \sin v - \delta y \cos v$ appearing in Q_- is zero if and only if the field point $x^a + \delta x^a$ lies in the beaming direction of the kink: see Appendix A.3.)

However, for the purposes of calculating the near-field derivatives $\partial_c h_{ab}$ it is necessary to retain the term $\eta_{ab}\delta x^a\delta x^b$ in full, with the result that many of the derivatives are undefined in the limit as $\delta x^a \rightarrow 0$. The near-field expansion of h_{ab} appearing in the footnote in Section 4.2 of Paper I is as complicated as it is because the “off-shell” contributions have been included so as to give a hint of the complexity of the derivatives.

It should be clear from the foregoing remarks that the breakdown of the field equations at the kinks is not strictly speaking a strong-field effect, but stems rather from the zero-thickness assumption underpinning the Nambu-Goto action, which imposes infinite worldsheet curvature at the kinks. It seems unlikely that the singularity at the kinks would be resolved in a fully non-linear solution of the Einstein equations, and I suspect that the kinks can be smoothed only by the inclusion of a finite-thickness field-theoretic source. However, the fact – mentioned in the footnote in Section 8 – that the Newtonian gravitational potential M_r/r remains bounded near the kinks (and in fact is likely to be of order μ there) offers some hope that the kinks on the ACO loop could be incorporated into a fully non-linear solution.

There are other topics of interest in this area that would repay further investigation. For example, it would be useful to be able to offer a definitive proof that the ACO loop does indeed have the lowest radiative efficiency of all flat-space cosmic string loop solutions, or to demonstrate that the evaporating ACO loop is at least neutrally stable to (cusp-free) small perturbations.

I would like to thank the referee for highlighting the need for some helpful clarifications.

A. Some Properties of the Roots V_1 and V_0

A.1. Bounds on V_1 and V_0

If the vector (X, Y, Z) corresponding to the spatial components $\bar{\mathbf{x}} = \frac{1}{4\pi} L_0(X, Y, Z)$ of the image point is represented in standard spherical polar coordinates:

$$(X, Y, Z) = (r \sin \theta \cos \phi, r \sin \theta \sin \phi, r \cos \theta), \quad (\text{A.1})$$

then since $T = r$ the equation (5.33) for the roots V_k specialized to V_1 reads

$$V_1 + \pi = r - [(r \sin \theta \cos \phi - \cos V_1)^2 + (r \sin \theta \sin \phi - \sin V_1)^2 + (r \cos \theta - \frac{1}{2}\pi)^2]^{1/2}. \quad (\text{A.2})$$

Differentiating the expression inside the square root with respect to V_1 indicates that $V_1 + \pi$ takes on an extremal value whenever $\sin(V_1 - \phi) = 0$ – unless $r \sin \theta = 0$, in which case equation (A.2) solves explicitly to give $V_1 + \pi = r - [1 + (r \mp \frac{1}{2}\pi)^2]^{1/2}$ (with the sign choice $\mp$ corresponding to $\cos \theta = \pm 1$ if $r \neq 0$). The case $r \sin \theta = 0$ is of course equivalent to $X = Y = 0$. and is discussed in more detail at the end of Section 5.2 It is easily verified that, in this case, $V_1 + \pi$ is a strictly monotonically increasing function of r , and so lies in the range $[-(1 + \frac{1}{4}\pi^2)^{1/2}, -\frac{1}{2}\pi]$ if $\cos \theta = +1$ and in $[-(1 + \frac{1}{4}\pi^2)^{1/2}, \frac{1}{2}\pi]$ if $\cos \theta = -1$.

If $\sin(V_1 - \phi) \neq 0$ then the right-hand side of (A.2) is extremized if either $(\cos V_1, \sin V_1) = (\cos \phi, \sin \phi)$ or $(\cos V_1, \sin V_1) = (-\cos \phi, -\sin \phi)$. It follows from (A.2) that

$$r - [(r \sin \theta + 1)^2 + (r \cos \theta - \frac{1}{2}\pi)^2]^{1/2} \leq V_1 + \pi \leq r - [(r \sin \theta - 1)^2 + (r \cos \theta - \frac{1}{2}\pi)^2]^{1/2}. \quad (\text{A.3})$$

Again, differentiation with respect to r of the lower and upper bounding functions in (A.3) is sufficient to show that both are monotonically increasing functions of r (and furthermore the left-hand function is strictly monotonic unless $\tan \theta = -2/\pi$,

while the right-hand function is strictly monotonic unless $\tan \theta = 2/\pi$). From this fact it is readily seen that

$$-(1 + \frac{1}{4}\pi^2)^{1/2} \leq V_1 + \pi \leq (1 + \frac{1}{4}\pi^2)^{1/2} \quad (\text{A.4})$$

(the lower bound being achieved when $r = 0$ or $\tan \theta = -2/\pi$, and the upper bound when $\tan \theta = 2/\pi$ and $r \geq (1 + \frac{1}{4}\pi^2)^{1/2}$). Since the root V_0 lies in the interval $[V_1, V_1 + 2\pi]$, it is evident that V_0 is also bounded:

$$-(1 + \frac{1}{4}\pi^2)^{1/2} - \pi \leq V_0 \leq (1 + \frac{1}{4}\pi^2)^{1/2} + \pi \approx 5.00369. \quad (\text{A.5})$$

The fact that V_0 and V_1 are bounded should come as no surprise, as V_0 and $V_1 + \pi$ are the values of the scaled time coordinate T at the points of intersection of the backwards light cone of the image point $[|\bar{\mathbf{x}}|, \bar{\mathbf{x}}]$ with the trajectories of the lower ($u = 0$) and upper ($u = \pm\pi$) kink points respectively.

A.2. The zeroes of $T - s(Z + \frac{1}{2}\pi) - v$

According to (5.34), the function $v(u)$ describing the curve of intersection Γ of the world sheet with the backwards light cone of $[|\bar{\mathbf{x}}|, \bar{\mathbf{x}}] \equiv \frac{1}{4\pi}L_0[T, X, Y, Z]$ has a possible stationary point whenever $E \equiv T - s(Z + \frac{1}{2}\pi) - v = 0$. (Recall here that $s \equiv \text{sgn}(u)$, or equivalently s is the sign of $\hat{\mathbf{z}} \cdot \mathbf{X}_u$ on the segment of the string loop at which the intersection point occurs.) It is important to be able to locate those points $[|\bar{\mathbf{x}}|, \bar{\mathbf{x}}]$ on the surface F_0 at which E goes to zero for some value of v in $[V_1, V_1 + 2\pi]$, because E appears as the denominator in the integrals contributing to $h_{ab}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}})$ in (5.36) and so the components of h_{ab} are potentially divergent at such a point.

As mentioned in section 5.2, E is positive semi-definite by virtue of equation (5.31), which can be rewritten in the form

$$\begin{aligned} T - s(Z + \frac{1}{2}\pi) - v &= u - s(Z + \frac{1}{2}\pi) + [(X - \cos v)^2 \\ &\quad + (Y - \sin v)^2 + \{Z - (|u| - \frac{1}{2}\pi)\}^2]^{1/2} \\ &\geq u - s(Z + \frac{1}{2}\pi) + |Z - (|u| - \frac{1}{2}\pi)|, \end{aligned} \quad (\text{A.6})$$

Here, equality occurs if and only if $X = \cos v$ and $Y = \sin v$, while the function on the third line is zero if and only if $u \leq s(Z + \frac{1}{2}\pi)$.

Suppose therefore that the point (X, Y) lies on the unit circle, so that $(X, Y) = (\cos v_F, \sin v_F)$ for some angle v_F . Then $T - s(Z + \frac{1}{2}\pi) - v_F$ will be zero whenever

$$Z = \lambda - \frac{1}{2}\pi \quad \text{and} \quad T = v_F + s\lambda, \quad (\text{A.7})$$

where λ is any number satisfying $\lambda \geq u$ if $u > 0$, or $\lambda \leq -u$ if $u < 0$. (Note that if the null constraint $T = (X^2 + Y^2 + Z^2)^{1/2}$ is also imposed then $v_F = [1 + (\lambda - \frac{1}{2}\pi)^2]^{1/2} - s\lambda$, and for each value of s there is a one-parameter family of spacetime points on F_0 for which $E = 0$ at some point on Γ .)

If $0 \leq \lambda \leq \pi$ then the spacetime point $\frac{1}{4\pi}L_0[T, X, Y, Z]$ coincides with the point $X^a(u, v) \equiv X^a(s\lambda, v_F)$ on the world sheet. It follows from the results of the previous paragraph that E will be zero for all values of u in the range $[0, \lambda]$ if $s = 1$, and all values of u in the range $[-\pi, -\lambda]$ if $s = -1$. This in turn means that the curve Γ includes a horizontal segment with $v \equiv v_F$ over the ranges of u indicated.

More important for the purposes of calculating $h_{ab}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}})$ are image points that lie off the worldsheet. These have $|Z| > \frac{1}{2}\pi$, and so either $\lambda > \pi$ if $s = 1$ or $\lambda < 0$ if $s = -1$. In geometric terms, such points either lie on the upwards ($Z > \frac{1}{2}\pi$) extension of the helical segment of the ACO loop with $u > 0$, or on the downwards ($Z < -\frac{1}{2}\pi$) extension of the helical segment with $u < 0$. Furthermore, it is evident from the preceding analysis that in this case E will be zero on the curve Γ for all values of $u \in [0, \pi]$ if $s = 1$, and all values of $u \in [-\pi, 0]$ if $s = -1$, and so Γ includes a straight-line segment with $v \equiv v_F$ over one of these ranges. An immediate consequence is that $V_0 = V_1 \equiv v_F$ if $s = 1$, and $V_0 = V_1 + 2\pi \equiv v_F$ if $s = -1$. (This straight-line segment corresponds to the “trivial solution” $V_k = v$ mentioned in Section 6.1.)

It is evident now that if the image point $[|\bar{\mathbf{x}}|, \bar{\mathbf{x}}]$ lies on one of the two helical extensions of the loop, but not on the loop itself, then the corresponding integral in the equation (5.36) for $h_{ab}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}})$ is undefined. However, the problematic integrals can be reformulated by making use of the equation (5.34) for dv/du to replace v with u as the variable of integration. Furthermore, since $(X, Y) = (\cos v_F, \sin v_F)$ the denominator $T - (u + v) + X \sin v - Y \cos v$ becomes simply $T - u - v_F$, where $v_F = \chi_{\pm} \equiv T \pm (Z + \frac{1}{2}\pi)$ if $s = \mp 1$. Hence,

$$\begin{aligned} \int_{V_0}^{V_1+2\pi} [T + (Z + \frac{1}{2}\pi) - v]^{-1} \bar{\Psi}_{ab}(-1, v) dv & \quad \text{is replaced by} \\ \int_{-\pi}^0 [-(Z + \frac{1}{2}\pi) - u]^{-1} \bar{\Psi}_{ab}(-1, \chi_+) du \end{aligned}$$

and

$$\int_{V_1}^{V_0} [T - (Z + \frac{1}{2}\pi) - v]^{-1} \bar{\Psi}_{ab}(+1, v) dv \quad \text{is replaced by}$$

$$\int_0^\pi (Z + \tfrac{1}{2}\pi - u)^{-1} \bar{\Psi}_{ab}(+1, \chi_-) du.$$

The new integrals are easily evaluated, giving

$$I_+ = -\ln[(\tfrac{1}{2}\pi - Z)/(-\tfrac{1}{2}\pi - Z)], \quad S_+ = I_+ \sin(\chi_+), \quad C_+ = I_+ \cos(\chi_+) \quad (\text{A.8})$$

(if $s = -1$) and

$$I_- = -\ln[(Z + \tfrac{1}{2}\pi)/(Z - \tfrac{1}{2}\pi)], \quad S_- = I_- \sin(\chi_-), \quad C_- = I_- \cos(\chi_-) \quad (\text{A.9})$$

(if $s = 1$) in place of the corresponding terms in (5.38)-(5.40).

A.3. The zeroes of $T - (u + v) + X \sin v - Y \cos v$

A second consequence of (5.34) is that the graph of v against u on the intersection curve Γ typically has a point of vertical inflection whenever $H \equiv T - (u + v) + X \sin v - Y \cos v = 0$. As was indicated in Section 5.2, the factor H is positive semi-definite, because

$$\begin{aligned} T - (u + v) + X \sin v - Y \cos v &= [(X \sin v - Y \cos v)^2 + (X \cos v + Y \sin v - 1)^2 \\ &\quad + \{Z - (|u| - \tfrac{1}{2}\pi)\}^2]^{1/2} + X \sin v - Y \cos v \\ &\geq |X \sin v - Y \cos v| + X \sin v - Y \cos v, \end{aligned} \quad (\text{A.10})$$

with equality occurring if and only if $X \cos v + Y \sin v = 1$ and $Z = |u| - \tfrac{1}{2}\pi$. In particular, it follows that spacetime points with $H = 0$ somewhere on Γ have $|Z| \leq \tfrac{1}{2}\pi$. It was seen earlier that spacetime points outside the world sheet with $E = 0$ somewhere on Γ have $|Z| > \tfrac{1}{2}\pi$, so the two classes of points are disjoint outside the world sheet.

Suppose now that $H = 0$ at the point $(u, v) = (u_F, v_F)$ on Γ . The equations $H = 0$ and $X \cos v + Y \sin v = 1$ (with $X \sin v - Y \cos v \leq 0$) can then be solved exactly to give

$$\begin{aligned} T &= u_F + v_F + (R^2 - 1)^{1/2}, & X &= \cos v_F - (R^2 - 1)^{1/2} \sin v_F \\ \text{and} \quad Y &= \sin v_F + (R^2 - 1)^{1/2} \cos v_F, \end{aligned} \quad (\text{A.11})$$

while of course $Z = |u_F| - \tfrac{1}{2}\pi$. The value of $R \geq 1$ is fixed by the null constraint $T = (X^2 + Y^2 + Z^2)^{1/2}$, which reads explicitly

$$u_F + v_F = [R^2 + (|u_F| - \tfrac{1}{2}\pi)^2]^{1/2} - (R^2 - 1)^{1/2}. \quad (\text{A.12})$$

So there exists a two-parameter family of spacetime points with the property that $H = 0$ somewhere on Γ .

If $[T, X, Y, Z]$ is held constant and u varied from u_F to $u_F + \delta u$, a perturbation expansion of the equation (5.28) for v indicates that

$$v(u_F + \delta u) - v_F = -(6\delta u)^{1/3} \quad (\text{A.13})$$

to leading order in δu . Hence, v remains a monotonic decreasing function of u even in the singular case where $H = 0$ somewhere on Γ .

The importance of the zeroes of H lies in the fact that the spacetime derivatives $\bar{\partial}_c h_{ab}$ of the metric perturbations $h_{ab}(\bar{t}, \bar{\mathbf{x}})$ are potentially divergent at any image point $[\bar{t}, \bar{\mathbf{x}}] \equiv \frac{1}{4\pi} L_0[T, X, Y, Z]$ with the property that $H = 0$ when $v = V_0$ or V_1 . This can be seen by first differentiating the equation (5.36) for h_{ab} and then integrating by parts to give

$$\begin{aligned} \bar{\partial}_c h_{ab} = & -4\mu(\bar{\partial}_c \chi_+) \int_{V_0}^{V_1+2\pi} (\chi_+ - v)^{-1} \Omega_{ab}(-1, v) dv \\ & -4\mu(\bar{\partial}_c \chi_-) \int_{V_1}^{V_0} (\chi_- - v)^{-1} \Omega_{ab}(+1, v) dv \\ & -4\mu(\chi_+ - V_1 - 2\pi)^{-1} \bar{\Psi}_{ab}(-1, V_1 + 2\pi) \bar{\partial}_c(V_1 - \chi_+) \\ & +4\mu(\chi_- - V_1)^{-1} \bar{\Psi}_{ab}(+1, V_1) \bar{\partial}_c(V_1 - \chi_-) \\ & -4\mu(\chi_- - V_0)^{-1} \bar{\Psi}_{ab}(+1, V_0) \bar{\partial}_c(V_0 - \chi_-) \\ & +4\mu(\chi_+ - V_0)^{-1} \bar{\Psi}_{ab}(-1, V_0) \bar{\partial}_c(V_0 - \chi_+), \end{aligned} \quad (\text{A.14})$$

where again $\chi_{\pm} = T \pm (Z + \frac{\pi}{2})$ and $\Omega_{ab}(s, v) = \partial_v \bar{\Psi}_{ab}(s, v)$ as defined in Section 6.1.

The two integrals on the right-hand side of (A.14) converge because all the components of Ω_{ab} are identical to components of $\bar{\Psi}_{ab}$, and it was demonstrated in the previous sub-section that integrals of this type will converge unless the field point lies on the worldsheet itself. Also, the derivatives $\bar{\partial}_c \chi_{\pm} = 4\pi L_0^{-1}[1, \pm \hat{\mathbf{z}}]_c$ are everywhere well defined.

However, the terms involving $\bar{\partial}_c V_1$ and $\bar{\partial}_c V_0$ in (A.14) are potentially problematic. Implicit differentiation of the equation (5.28) defining v gives:

$$\begin{aligned} \bar{\partial}_c v = & [T - (u + v) + X \sin v - Y \cos v]^{-1} \\ & \times 4\pi L_0^{-1}[T - (u + v), -(X - \cos v), -(Y - \sin v), -\{Z - (|u| - \frac{1}{2}\pi)\}]_c, \end{aligned} \quad (\text{A.15})$$

where the denominator is of course just H . The derivative $\bar{\partial}_c V_k$ therefore typically diverges if $H = 0$ at $v = V_k$ (and in fact the components of the 4-vector multiplying H^{-1} in (A.15) will all vanish only if the point $\frac{1}{4\pi}L_0[T, X, Y, Z]$ lies on the world sheet).¹¹

In fact, if $k = 0$ or 1 and the image point $\bar{x}^a = \frac{1}{4\pi}L_0[T, X, Y, Z]^a$ has the property that

$$X \cos V_k + Y \sin V_k = 1, \quad T = k\pi + V_k - X \sin V_k + Y \cos V_k \quad \text{and} \quad Z = (|k| - \frac{1}{2})\pi \quad (\text{A.16})$$

(with $X \sin V_k - Y \cos V_k \leq 0$) then the perturbative solution of the equation (5.33) for V_k at a nearby point $\bar{x}^a + \delta\bar{x}^a = \frac{1}{4\pi}L_0[T + \delta T, X + \delta X, Y + \delta Y, Z + \delta Z]^a$ reads:

$$\begin{aligned} V_k(\bar{x}^a + \delta\bar{x}^a) - V_k(\bar{x}^a) &= 6^{1/3}(\delta T + \delta X \sin V_k - \delta Y \cos V_k)^{1/3} \\ &\quad + \frac{1}{4}(X \sin V_k - Y \cos V_k)^{-1} 6^{2/3}(\delta T + \delta X \sin V_k - \delta Y \cos V_k)^{2/3}, \end{aligned} \quad (\text{A.17})$$

plus terms linear in $\delta\bar{x}^a$.

Substitution of the expansion (A.17) into the equation (5.36) for h_{ab} then gives:

$$h_{tz}(\bar{x}^a + \delta\bar{x}^a) - h_{tz}(\bar{x}^a) = -8\hat{z}\mu(\delta_k^{1/3} + \frac{1}{4}\delta_k^{2/3}) + O(\delta\bar{x}), \quad (\text{A.18})$$

$$h_{xz}(\bar{x}^a + \delta\bar{x}^a) - h_{xz}(\bar{x}^a) = -8\hat{z}\mu[\delta_k^{1/3} \sin V_k + \frac{1}{4}\delta_k^{2/3}(\sin V_k + 2D_k \cos V_k)] + O(\delta\bar{x}) \quad (\text{A.19})$$

and

$$h_{yz}(\bar{x}^a + \delta\bar{x}^a) - h_{yz}(\bar{x}^a) = 8\hat{z}\mu[\delta_k^{1/3} \cos V_k + \frac{1}{4}\delta_k^{2/3}(\cos V_k - 2D_k \sin V_k)] + O(\delta\bar{x}), \quad (\text{A.20})$$

where

$$\delta_k \equiv 6D_k^{-3}(\delta T + \delta X \sin V_k - \delta Y \cos V_k), \quad (\text{A.21})$$

¹¹It might be supposed from (A.14) that $\bar{\partial}_c h_{ab}$ is also undefined whenever $\chi_{\pm} - V = 0$, with $V = V_1, V_0$ or V_{-1} . That this is not so (at least for image points outside the world sheet) can be seen by replacing v with u as the integration variable in the equation (5.36) for h_{ab} to give:

$$h_{ab}(|\bar{\mathbf{x}}|, \bar{\mathbf{x}}) = -4\mu \int_{-\pi}^0 H^{-1} \bar{\Psi}_{ab}(-1, v) du - 4\mu \int_0^{\pi} H^{-1} \bar{\Psi}_{ab}(+1, v) dv$$

It is clear that $\bar{\partial}_c h_{ab}$ involves only terms of the form $\Omega_{ab} \bar{\partial}_c v$ and $H^{-2} \bar{\partial}_c H$, where $\bar{\partial}_c H$ is a linear combination of $\bar{\partial}_c T$, $\bar{\partial}_c X$, $\bar{\partial}_c Y$ and $\bar{\partial}_c v$, and according to (A.15) $\bar{\partial}_c v$ is undefined only when $H = 0$.

$$D_k \equiv -(X \sin V_k - Y \cos V_k) \geq 0 \quad (\text{A.22})$$

and $\hat{z} \equiv \text{sgn}(Z) \equiv \text{sgn}(k - \frac{1}{2})$, while all other components of $h_{ab}(\bar{x}^a + \delta\bar{x}^a) - h_{ab}(\bar{x}^a)$ are of linear order or higher in the components of $\delta\bar{x}$.¹²

In summary, if the field point $\bar{x}^a$ satisfies the constraints (A.16) for either $k = 0$ or $k = 1$ then the derivatives of the metric components h_{tz} , h_{xz} and h_{yz} with respect to $\bar{t}$, $\bar{x}$ and $\bar{y}$ are all undefined there. Field points of this type have $\bar{z} = \pm \frac{1}{8}L_0$ and so have the same altitude as one of the kink points on the string. The failure of h_{ab} to be differentiable at such points therefore has no effect on the analysis of the first-order string equation of motion in Section 6, as the equation of motion is there developed and solved only for points on the string away from the kinks. However, the singularities in the derivatives of h_{ab} do potentially affect the matching of the weak-field solution to the Minkowski vacuum across the null surface F_L . Fortunately, the relevant combinations of the components of h_{ab} are all differentiable on F_L , as is shown in detail in Section 7.2.

It might seem puzzling that the weak-field metric induced by the ACO loop is singular on a manifold of points that extends away from the worldsheet. However, this pathological behavior has a simple physical explanation. It is well known ([23], Section 6.3) that a generic kink on a zero-thickness string emits a narrow beam of gravitational radiation in the direction of its motion. Although the metric derivative is singular on the locus of the beam, the singularity is nonetheless relatively benign, as the geodesic equations for a free particle remain integrable across the beam.

In the case of the ACO loop, the kink points correspond to $u = k\pi$ for $k = 0$ or 1 and so trace out the two families of spacetime points

$$X_{(0)}^a = \frac{1}{4\pi}L_0[k\pi + v, \cos v, \sin v, (|k| - \frac{1}{2})\pi]^a. \quad (\text{A.23})$$

The instantaneous 4-velocities of the kink points are therefore

$$\partial_v X_{(0)}^a = \frac{1}{4\pi}L_0[1, -\sin v, \cos v, 0]^a, \quad (\text{A.24})$$

¹²Incidentally, the factor D_k^{-3} appearing in the equation for δ_k is guaranteed to be well defined if the field point lies away from the string worldsheet, because if $X \sin V_k - Y \cos V_k = 0$ then the previous constraints (A.16) imply that

$$X = \cos V_k, \quad Y = \sin V_k, \quad T = k\pi + V_k \quad \text{and} \quad Z = |k\pi| - \frac{1}{2}\pi$$

and so the field point is identical to the point $(u, v) = (k\pi, V_k)$ on the worldsheet.

and the beams, which trace out null curves with tangent vector $\partial_v X_{(0)}^a$, can be represented parametrically in the form

$$[T, X, Y, Z] = [k\pi + v, \cos v, \sin v, (|k| - \frac{1}{2})\pi] + \lambda[1, -\sin v, \cos v, 0], \quad (\text{A.25})$$

with λ an affine parameter. Hence $Z = (|k| - \frac{1}{2})\pi$ and, after eliminating λ ,

$$T - (k\pi + v) + X \sin v - Y \cos v = 0. \quad (\text{A.26})$$

The last equation is just the constraint $H = 0$ with $u = k\pi$. So a spacetime point lying off the worldsheet satisfies the constraints (A.16) if and only if it lies in the beaming direction of a kink.

B. Calculating h_{ab} and its Derivatives

With $\chi_{\pm} \equiv v + (u \pm u) + 4\pi L_0^{-1}(\delta t \pm \delta z)$ and

$$\begin{aligned} \chi_+ - V_0 &= 2u + 8\pi L_0^{-1}\delta z, & \chi_+ - V_1 &= W + 2u + 4\pi L_0^{-1}(\delta t + \delta z - \delta V) \\ \chi_- - V_0 &= \frac{1}{2}u^{-1}(4\pi L_0^{-1})^2 Q_+ & \text{and} & \quad \chi_- - V_1 = W + 4\pi L_0^{-1}(\delta t - \delta z - \delta V), \end{aligned} \quad (\text{B.1})$$

where δV is of linear order in δx^a and is defined in (6.13), while Q_+ is of quadratic order and is defined in (6.15), the expressions (5.38)-(5.40) become, for $u > 0$ and δx^a small,

$$I_+ = \ln(W + 2u - 2\pi) - \ln(2u) + 4\pi L_0^{-1} \left(\frac{\delta t + \delta z - \delta V}{W + 2u - 2\pi} - \frac{\delta z}{u} \right), \quad (\text{B.2})$$

$$I_- = \ln[(4\pi L_0^{-1})^2 Q_+] - \ln(2uW) - 4\pi L_0^{-1} \left(\frac{\delta t - \delta z - \delta V}{W} \right), \quad (\text{B.3})$$

$$\begin{aligned} \begin{bmatrix} S_+ \\ C_+ \end{bmatrix} &= \begin{bmatrix} -\cos(2u + v) & \sin(2u + v) \\ \sin(2u + v) & \cos(2u + v) \end{bmatrix} \begin{bmatrix} \text{Si}(W + 2u - 2\pi) - \text{Si}(2u) \\ \text{Ci}(W + 2u - 2\pi) - \text{Ci}(2u) \end{bmatrix} \\ &\quad + 4\pi L_0^{-1} \begin{bmatrix} \sin(2u + v) & \cos(2u + v) \\ \cos(2u + v) & -\sin(2u + v) \end{bmatrix} \begin{bmatrix} \text{Si}(W + 2u - 2\pi) - \text{Si}(2u) \\ \text{Ci}(W + 2u - 2\pi) - \text{Ci}(2u) \end{bmatrix} (\delta t + \delta z) \\ &\quad - 4\pi L_0^{-1} u^{-1} \begin{bmatrix} \sin v \\ \cos v \end{bmatrix} \delta z + 4\pi L_0^{-1} (W + 2u - 2\pi)^{-1} \begin{bmatrix} \sin(v - W) \\ \cos(v - W) \end{bmatrix} (\delta t + \delta z - \delta V) \end{aligned} \quad (\text{B.4})$$

and

$$\begin{aligned}
\begin{bmatrix} S_- \\ C_- \end{bmatrix} &= \{ \ln[(4\pi L_0^{-1})^2 Q_+] + \gamma_E - \ln(2u) \} \begin{bmatrix} \sin v \\ \cos v \end{bmatrix} - \begin{bmatrix} -\cos v & \sin v \\ \sin v & \cos v \end{bmatrix} \begin{bmatrix} \text{Si } W \\ \text{Ci } W \end{bmatrix} \\
&+ 4\pi L_0^{-1} \{ \ln[(4\pi L_0^{-1})^2 Q_+] + \gamma_E - \ln(2u) \} \begin{bmatrix} \cos v \\ -\sin v \end{bmatrix} (\delta t - \delta z) \\
&- 4\pi L_0^{-1} \begin{bmatrix} \sin v & \cos v \\ \cos v & -\sin v \end{bmatrix} \begin{bmatrix} \text{Si } W \\ \text{Ci } W \end{bmatrix} (\delta t - \delta z) \\
&- 4\pi L_0^{-1} W^{-1} \begin{bmatrix} \sin(v - W) \\ \cos(v - W) \end{bmatrix} (\delta t - \delta z - \delta V). \tag{B.5}
\end{aligned}$$

Note here that $\text{Ci } x = \ln x + \gamma_E + O(x^2)$ for small values of x , where $\gamma_E \approx 0.5772$ is Euler's constant, while $\text{Si } x = x + O(x^3)$.

If the terms which tend to zero as $\delta x^a \rightarrow 0$ in (B.2)-(B.5) are neglected, and the resulting leading-order expansions substituted into (5.37), the metric perturbations h_{ab} take on the form shown in equation (6.16).

Also, given that

$$\begin{aligned}
\partial(\delta V)/\partial(\delta x^a) &= (u - \pi + W - \sin W)^{-1} \\
&\times [u - \pi + W, \{ \cos(v - W) - \cos v \} \hat{\mathbf{x}} + \{ \sin(v - W) - \sin v \} \hat{\mathbf{y}} + (\pi - u) \hat{\mathbf{z}}]_a \tag{B.6}
\end{aligned}$$

the terms in the small-distance expansions of the derivatives of $I_\pm$, $S_\pm$ and $C_\pm$ which do not vanish in the limit as $\delta x^a \rightarrow 0$ can be read off directly from equations (B.2)-(B.5):

$$\partial_a I_+ = 4\pi L_0^{-1} \frac{1}{(W+2u-2\pi)(u-\pi+W-\sin W)} D_a - 4\pi L_0^{-1} u^{-1} [0, \hat{\mathbf{z}}]_a \tag{B.7}$$

$$\partial_a I_- = \partial_a \Lambda + 4\pi L_0^{-1} \frac{1}{W(u-\pi+W-\sin W)} E_a \tag{B.8}$$

$$\begin{aligned}
\begin{bmatrix} \partial_a S_+ \\ \partial_a C_+ \end{bmatrix} &= 4\pi L_0^{-1} \begin{bmatrix} \sin(2u+v) & \cos(2u+v) \\ \cos(2u+v) & -\sin(2u+v) \end{bmatrix} \begin{bmatrix} \text{Si } (W+2u-2\pi) - \text{Si } (2u) \\ \text{Ci } (W+2u-2\pi) - \text{Ci } (2u) \end{bmatrix} [1, \hat{\mathbf{z}}]_a \\
&- 4\pi L_0^{-1} u^{-1} \begin{bmatrix} \sin v \\ \cos v \end{bmatrix} [0, \hat{\mathbf{z}}]_a + 4\pi L_0^{-1} \frac{1}{(W+2u-2\pi)(u-\pi+W-\sin W)} \begin{bmatrix} \sin(v-W) \\ \cos(v-W) \end{bmatrix} D_a \tag{B.9}
\end{aligned}$$

and

$$\begin{aligned}
\begin{bmatrix} \partial_a S_- \\ \partial_a C_- \end{bmatrix} &= \partial_a \Lambda \begin{bmatrix} \sin v \\ \cos v \end{bmatrix} + 4\pi L_0^{-1} \{ \Lambda + \gamma_E - \ln(2u) \} \begin{bmatrix} \cos v \\ -\sin v \end{bmatrix} [1, -\hat{\mathbf{z}}]_a \\
&+ 4\pi L_0^{-1} \partial_a \Lambda \begin{bmatrix} \cos v \\ -\sin v \end{bmatrix} (\delta t - \delta z) - 4\pi L_0^{-1} \begin{bmatrix} \sin v & \cos v \\ \cos v & -\sin v \end{bmatrix} \begin{bmatrix} \text{Si } W \\ \text{Ci } W \end{bmatrix} [1, -\hat{\mathbf{z}}]_a \\
&+ 4\pi L_0^{-1} \frac{1}{W(u-\pi+W-\sin W)} \begin{bmatrix} \sin(v-W) \\ \cos(v-W) \end{bmatrix} E_a.
\end{aligned} \tag{B.10}$$

where $\Lambda = \ln[(4\pi L_0^{-1})^2 Q_+]$,

$$D_a \equiv [-\sin W, -\{\cos(v-W) - \cos v\} \hat{\mathbf{x}} - \{\sin(v-W) - \sin v\} \hat{\mathbf{y}} + (2u - 2\pi + W - \sin W) \hat{\mathbf{z}}]_a \tag{B.11}$$

and

$$E_a \equiv [\sin W, \{\cos(v-W) - \cos v\} \hat{\mathbf{x}} + \{\sin(v-W) - \sin v\} \hat{\mathbf{y}} + (W - \sin W) \hat{\mathbf{z}}]_a. \tag{B.12}$$

Again, the expressions (6.21)-(6.24) for the spacetime derivatives of h_{ab} at points on the null surface F_0 and close to the worldsheet of the loop follow directly by substituting the leading-order expansions (B.7)-(B.10) into (5.37).

C. Deriving the Equations of Motion

If the position function X^a of the evaporating loop is assumed to have the form (4.20) then

$$X_{(0)}^a(u, v) = \frac{1}{4\pi} L_0 [u + v, \mathbf{r}(u, v)]^a \tag{C.1}$$

with

$$\mathbf{r}(u, v) = \cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}} + (|u| - \frac{1}{2}\pi) \hat{\mathbf{z}}, \tag{C.2}$$

and the first-order perturbation δX^a is as given in (4.23), namely

$$\delta X^a(u, v) = -\frac{1}{4\pi} L_0 \kappa \mu \left[\frac{1}{2}(u+v)^2, (u+v) \mathbf{r}(u, v) \right]^a + \frac{1}{4\pi} L_0 \mu [-F, G\{\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}\} + A \hat{\mathbf{z}}]^a \tag{C.3}$$

The first derivatives of $X_{(0)}^a$ are therefore:

$$X_{(0),u}^a = \frac{1}{4\pi} L_0 [1, s \hat{\mathbf{z}}]^a \quad \text{and} \quad X_{(0),v}^a = \frac{1}{4\pi} L_0 [1, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}}]^a, \tag{C.4}$$

with $s = \text{sgn}(u)$, while the only non-zero component of $X_{(0),AB}^a$ is¹³:

$$X_{(0),vv}^a = \frac{1}{4\pi} L_0 [0, -\cos(v) \hat{\mathbf{x}} - \sin(v) \hat{\mathbf{y}}]^a. \quad (\text{C.5})$$

Similarly, the first derivatives of δX^a are:

$$\delta X^a_{,u} = -\frac{1}{4\pi} L_0 \kappa \mu [u+v, \mathbf{r}(u, v) + s(u+v) \hat{\mathbf{z}}]^a + \frac{1}{4\pi} L_0 \mu [-F_u, G_u \{\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}\} + A_u \hat{\mathbf{z}}]^a, \quad (\text{C.6})$$

$$\begin{aligned} \delta X^a_{,v} = & -\frac{1}{4\pi} L_0 \kappa \mu [u+v, \mathbf{r}(u, v)]^a + \frac{1}{4\pi} L_0 \mu [-F_v, G_v \{\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}\} + A_v \hat{\mathbf{z}}]^a \\ & + \frac{1}{4\pi} L_0 \mu \{G - \kappa(u+v)\} [0, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}}]^a \end{aligned} \quad (\text{C.7})$$

and the only second derivative of δX^a contributing to the first-order equation of motion is:

$$\begin{aligned} \delta X^a_{,uv} = & -\frac{1}{4\pi} L_0 \kappa \mu [1, s \hat{\mathbf{z}}]^a + \frac{1}{4\pi} L_0 \mu (G_u - \kappa) [0, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}}]^a \\ & + \frac{1}{4\pi} L_0 \mu [-F_{uv}, G_{uv} \{\cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}\} + A_{uv} \hat{\mathbf{z}}]^a. \end{aligned} \quad (\text{C.8})$$

The 4-dimensional tangent space at the point $x^a = X^a(u, v)$ is spanned by the vectors $X_{(0),u}^a$ and $X_{(0),v}^a$, plus

$$M^a(s, v) = [1, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}} + s \hat{\mathbf{z}}]^a \quad \text{and} \quad N^a(v) = [0, \cos(v) \hat{\mathbf{x}} + \sin(v) \hat{\mathbf{y}}]^a. \quad (\text{C.9})$$

In calculating the various contributions to the first-order equation of motion (2.14), it is useful to note that all the terms are transvected with q_d^c , where $q_{ab} = -\bar{\Psi}_{ab}(s, v)$ is the projection operator orthogonal to the worldsheet at $X^a(u, v)$, so that

$$q_d^c X_{(0),u}^d = q_d^c X_{(0),v}^d = 0, \quad q_d^c M^d = M^c \quad \text{and} \quad q_d^c N^d = N^c. \quad (\text{C.10})$$

Other helpful identities involving q_d^c include:

$$q_d^c [0, -\sin(v) \hat{\mathbf{x}} + \cos(v) \hat{\mathbf{y}}]^d = M^c, \quad q_d^c [1, \mathbf{0}]^d = -M^c \quad \text{and} \quad q_d^c [0, s \hat{\mathbf{z}}]^d = M^c. \quad (\text{C.11})$$

¹³The claim that $X_{(0),uu}^a = 0$ is true away from the kink points at $u = 0$ and $u = \pi$, but it can be seen from the expression for $X_{(0),u}^a$ that the z -component of $X_{(0),uu}^a$ contains a distributional term at the kinks. The possibility of pathological behavior at the kinks will be ignored here, as the ramifications of the singularities at the kinks have already been discussed in Paper I. It seems unlikely that the dynamical role of the kinks can be clarified without developing a solution of the Abelian Higgs field equations as a finite-thickness source for the ACO loop.

Thus, recalling that the only non-zero component of γ^{AB} (at zeroth order in μ) is $\gamma^{uv} = (\frac{1}{4\pi}L_0)^{-2}$, the first term in (2.14) is:

$$\begin{aligned} q_d^c \gamma^{CD} \delta X^d{}_{,CD} &= 2(\frac{1}{4\pi}L_0)^{-2} q_d^c \delta X^d{}_{,uv} \\ &= 2\mu(\frac{1}{4\pi}L_0)^{-1} (F_{uv} + sA_{uv} + G_u - \kappa) M^c + 2\mu(\frac{1}{4\pi}L_0)^{-1} G_{uv} N^c. \end{aligned} \quad (\text{C.12})$$

The second term in (2.14) is identically zero by virtue of the zeroth-order equation of motion, while the third term is:

$$\begin{aligned} -2q_d^c \gamma^{AC} \gamma^{BD} \eta_{ab} X_{(0),A}^a \delta X^b{}_{,B} X_{(0),CD}^d &= -2(\frac{1}{4\pi}L_0)^{-4} q_d^c X_{(0),vv}^d (\eta_{ab} X_{(0),u}^a \delta X^b{}_{,u}) \\ &= 2\mu(\frac{1}{4\pi}L_0)^{-1} [\kappa s(|u| - \frac{1}{2}\pi) - (F_u + sA_u)] N^c. \end{aligned} \quad (\text{C.13})$$

The remaining terms in (2.14) are more complicated, as they involve h_{ab} or its derivatives. However, evaluation of these terms is simplified by making use of the following table, which represents the inner products of the terms in the first column with those in the first row in the form (k_1, k_2, k_3, k_4) , which in turn is shorthand for $k_1 X_{(0)a,u} + k_2 X_{(0)a,v} + k_3 M_a + k_4 N_a$. Also, c_2 and s_2 stand in for $\cos 2u$ and $\sin 2u$ respectively, while c_W and s_W are shorthand for $\cos W$ and $\sin W$.

	$X_{(0),u}^b$	$X_{(0),v}^b$
$\Psi_{ab}(s, v)$	$\mathbf{0}$	$\mathbf{0}$
$\Psi_{ab}(-s, v)$	$(0, 4, -2, 0)$	$\mathbf{0}$
$\Phi_{ab}(s, v)$	$\mathbf{0}$	$(-2, 0, 1, 0)$
$\Phi_{ab}(-s, 2u + v)$	$(-2, 0, 2, 0)c_2 + (0, 0, 0, -2)s_2$	$(-2, -2, 3, 0)c_2 + (0, 0, 0, -1)s_2$
$\Omega_{ab}(s, v)$	$\mathbf{0}$	$(0, 0, 0, -1)$
$\Omega_{ab}(-s, 2u + v)$	$(2, 0, -2, 0)s_2 + (0, 0, 0, -2)c_2$	$(2, 2, -3, 0)s_2 + (0, 0, 0, -1)c_2$
$\Pi_{ab}(s)$	$\mathbf{0}$	$(2, 0, -1, 0)$
$\Pi_{ab}(-s)$	$(2, 4, -2, 0)$	$(2, 2, -3, 0)$
$\Psi_{ab}(s, v - W)$	$\mathbf{0}$	$(2, 0, -1, 0)(1 - c_W) + (0, 0, 0, 1)s_W$
$\Psi_{ab}(-s, v - W)$	$(2(1 - c_W), 4, -2(2 - c_W), 2s_W)$	$(2, 2, -3, 0)(1 - c_W) + (0, 0, 0, 1)s_W$

Furthermore, the vectors $X_{(0),u}^a$ and $X_{(0),v}^a$ are both null and orthogonal to M^a and N^a , with $X_{(0),u} \cdot X_{(0),v} = (\frac{1}{4\pi}L_0)^2$.

It should of course be remembered that the expansions (6.16) and (6.21)-(6.24) were developed for the case $u > 0$ only, and so $s = +1$ in the expressions

in this table. Because the term involving $\ln[(4\pi L_0^{-1})^2 Q_+]$ in the equation for h_{ab} is proportional to $\bar{\Psi}_{ab}(+1, v)$, the term containing $h_{ab} X_{(0),A}^a X_{(0),B}^b$ in (2.14) is free of logarithmic singularities. Similarly, the singular terms proportional to $\bar{\Psi}_{ab}(+1, v)$ in (6.21)-(6.24) do not contribute to the term $X_{(0),B}^b (h_{bd,a} - \frac{1}{2} h_{ab,d})$ on the right-hand side of (2.14). However, there remain singular terms proportional to $\Omega_{ab}(+1, v)$ in the derivatives of h_{ab} whose contributions to the first-order equation of motion need to be examined more carefully.

Now, the fourth term in the equation of motion is:

$$-q_d^c \gamma^{AC} \gamma^{BD} h_{ab} X_{(0),A}^a X_{(0),B}^b X_{(0),CD}^d = -(\frac{1}{4\pi} L_0)^{-4} q_d^c X_{(0),vv}^d (h_{ab} X_{(0),u}^a X_{(0),u}^b) \quad (\text{C.14})$$

where the only non-zero term in $h_{ab} X_{(0),u}^a X_{(0),u}^b$ is proportional to the coefficient of $\Pi_{ab}(-1)$:

$$h_{ab} X_{(0),u}^a X_{(0),u}^b = 16\mu(\frac{1}{4\pi} L_0)^2 [\ln(W + 2u - 2\pi) - \ln(2u)], \quad (\text{C.15})$$

and so

$$-q_d^c \gamma^{AC} \gamma^{BD} h_{ab} X_{(0),A}^a X_{(0),B}^b X_{(0),CD}^d = 16\mu(\frac{1}{4\pi} L_0)^{-1} [\ln(W + 2u - 2\pi) - \ln(2u)] N^c. \quad (\text{C.16})$$

Furthermore, the right-hand expression in (2.14) can be broken into two parts:

$$-q^{cd} \gamma^{AB} X_{(0),A}^a X_{(0),B}^b h_{bd,a} = -(\frac{1}{4\pi} L_0)^{-2} q^{cd} (X_{(0),u}^a X_{(0),v}^b + X_{(0),v}^a X_{(0),u}^b) h_{bd,a} \quad (\text{C.17})$$

and

$$\frac{1}{2} q^{cd} \gamma^{AB} X_{(0),A}^a X_{(0),B}^b h_{ab,d} = (\frac{1}{4\pi} L_0)^{-2} q^{cd} X_{(0),u}^a X_{(0),v}^b h_{ab,d}. \quad (\text{C.18})$$

Here, $q^{cd} X_{(0),u}^a X_{(0),v}^b h_{bd,a}$ involves terms of the form

$$\begin{aligned} -q^{cd} X_{(0),v}^b \partial_t h_{bd} &= 4\mu(\text{Si } W) M^c + 4\mu(\delta t - \delta z) (\partial_t \Lambda) N^c + 4\mu[\Lambda + \gamma_E - \text{Ci } W - \ln(2u)] N^c \\ &\quad + 4\mu\Sigma(u) [-(3 \cos 2u) M^c + (\sin 2u) N^c] + 4\mu K(u) [(3 \sin 2u) M^c + (\cos 2u) N^c] \\ &\quad + 4\mu \frac{\sin W}{u-\pi+W-\sin W} \frac{1}{W} [(1 - \cos W) M^c - (\sin W) N^c] \\ &\quad - 4\mu \frac{\sin W}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} [3(1 - \cos W) M^c - (\sin W) N^c] \end{aligned} \quad (\text{C.19})$$

and

$$\begin{aligned} -q^{cd} X_{(0),v}^b \partial_z h_{bd} &= -4\mu(\text{Si } W) M^c + 4\mu(\delta t - \delta z) (\partial_z \Lambda) N^c - 4\mu[\Lambda + \gamma_E - \text{Ci } W - \ln(2u)] N^c \\ &\quad + 4\mu\Sigma(u) [-(3 \cos 2u) M^c + (\sin 2u) N^c] + 4\mu K(u) [(3 \sin 2u) M^c + (\cos 2u) N^c] \\ &\quad + 4\mu \frac{W - \sin W}{u-\pi+W-\sin W} \frac{1}{W} [(1 - \cos W) M^c - (\sin W) N^c] \\ &\quad + 4\mu \frac{2u-2\pi+W-\sin W}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} [3(1 - \cos W) M^c - (\sin W) N^c]. \end{aligned} \quad (\text{C.20})$$

So

$$\begin{aligned}
-\left(\frac{1}{4\pi}L_0\right)^{-2}q^{cd}X_{(0),u}^a X_{(0),v}^b h_{bd,a} &= -\left(\frac{1}{4\pi}L_0\right)^{-1}q^{cd}X_{(0),v}^b (\partial_t h_{bd} + \partial_z h_{bd}) \\
&= 4\mu\left(\frac{1}{4\pi}L_0\right)^{-1}\{(\delta t - \delta z)[(\partial_t \Lambda) + (\partial_z \Lambda)]N^c \\
&\quad + 2\Sigma(u)[-(3\cos 2u)M^c + (\sin 2u)N^c] \\
&\quad + 2K(u)[(3\sin 2u)M^c + (\cos 2u)N^c] \\
&\quad + \frac{1}{u-\pi+W-\sin W}[(1-\cos W)M^c - (\sin W)N^c] \\
&\quad + \frac{2u-2\pi+W-2\sin W}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} [3(1-\cos W)M^c - (\sin W)N^c]\},
\end{aligned} \tag{C.21}$$

where

$$\begin{aligned}
(\partial_t \Lambda) + (\partial_z \Lambda) &= Q_+^{-1} \partial_t (\bar{\Psi}_{ab} \delta x^a \delta x^b) + Q_+^{-1} \partial_z (\bar{\Psi}_{ab} \delta x^a \delta x^b) \\
&= 2Q_+^{-1} (\bar{\Psi}_{at} + \bar{\Psi}_{az}) \delta x^a = 0,
\end{aligned} \tag{C.22}$$

with $\bar{\Psi}_{ab}$ here denoting $\bar{\Psi}_{ab}(+1, v)$.

Similarly, $q^{cd}X_{(0),v}^a X_{(0),u}^b h_{bd,a}$ is a sum of three terms proportional to

$$\begin{aligned}
-q^{cd}X_{(0),u}^b \partial_t h_{bd} &= 4\mu\Sigma(u)[-(2\cos 2u)M^c + (2\sin 2u)N^c] + 4\mu K(u)[(2\sin 2u)M^c + (2\cos 2u)N^c] \\
&\quad - 8\mu \frac{\sin W}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} [(2-\cos W)M^c - (\sin W)N^c],
\end{aligned} \tag{C.23}$$

$$-q^{cd}X_{(0),u}^b \partial_x h_{bd} = -8\mu \frac{\cos(v-W)-\cos v}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} [(2-\cos W)M^c - (\sin W)N^c] \tag{C.24}$$

and

$$-q^{cd}X_{(0),u}^b \partial_y h_{bd} = -8\mu \frac{\sin(v-W)-\sin v}{u-\pi+W-\sin W} \frac{1}{W+2u-2\pi} [(2-\cos W)M^c - (\sin W)N^c]. \tag{C.25}$$

So

$$\begin{aligned}
-\left(\frac{1}{4\pi}L_0\right)^{-2}q^{cd}X_{(0),v}^a X_{(0),u}^b h_{bd,a} &= -\left(\frac{1}{4\pi}L_0\right)^{-1}q^{cd}X_{(0),u}^b [\partial_t h_{bd} - (\sin v)\partial_x h_{bd} + (\cos v)\partial_y h_{bd}] \\
&= 4\mu\left(\frac{1}{4\pi}L_0\right)^{-1}\Sigma(u)[-(2\cos 2u)M^c + (2\sin 2u)N^c] \\
&\quad + 4\mu\left(\frac{1}{4\pi}L_0\right)^{-1}K(u)[(2\sin 2u)M^c + (2\cos 2u)N^c].
\end{aligned} \tag{C.26}$$

Adding together (C.21) and (C.26) gives equation (6.30).

The various terms contributing to the second part of the last term in (2.14) are:

$$X_{(0),u}^a X_{(0),v}^b \partial_t h_{ab} = 8\mu\left(\frac{1}{4\pi}L_0\right)[K(u)\sin 2u - \Sigma(u)\cos 2u] - 8\mu\left(\frac{1}{4\pi}L_0\right) \frac{\sin W}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi}, \tag{C.27}$$

$$X_{(0),u}^a X_{(0),v}^b \partial_x h_{ab} = -8\mu \left(\frac{1}{4\pi} L_0\right) \frac{\cos(v-W) - \cos v}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi}, \quad (\text{C.28})$$

$$X_{(0),u}^a X_{(0),v}^b \partial_y h_{ab} = -8\mu \left(\frac{1}{4\pi} L_0\right) \frac{\sin(v-W) - \sin v}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi} \quad (\text{C.29})$$

and

$$X_{(0),u}^a X_{(0),v}^b \partial_z h_{ab} = 8\mu \left(\frac{1}{4\pi} L_0\right) [K(u) \sin 2u - \Sigma(u) \cos 2u] + 8\mu \left(\frac{1}{4\pi} L_0\right) \frac{2u-2\pi+W-\sin W}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi}. \quad (\text{C.30})$$

Now, if V_d is any contravariant vector and $s = +1$, then

$$q^{cd} V_d = -[V_t - (\sin v) V_x + (\cos v) V_y + V_z] M^c - [(\cos v) V_x + (\sin v) V_y] N^c. \quad (\text{C.31})$$

Hence, the second part of the last term in (2.14) becomes

$$\begin{aligned} \left(\frac{1}{4\pi} L_0\right)^{-2} q^{cd} X_{(0),u}^a X_{(0),v}^b h_{ab,d} &= -16\mu \left(\frac{1}{4\pi} L_0\right)^{-1} [K(u) \sin 2u - \Sigma(u) \cos 2u] M^c \\ &\quad - 8\mu \left(\frac{1}{4\pi} L_0\right)^{-1} \frac{W+2u-2\pi-\sin W}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi} M^c \\ &\quad - 8\mu \left(\frac{1}{4\pi} L_0\right)^{-1} \frac{1-\cos W}{u-\pi+W-\sin W} \frac{1-\cos W}{W+2u-2\pi} N^c. \end{aligned} \quad (\text{C.32})$$

D. The Intrinsic 2-Metric and the Length of the Evaporating ACO Loop

D.1. The intrinsic 2-metric γ_{AB}

To linear order in μ , the intrinsic 2-metric $\gamma_{AB} = g_{ab} X^a_{(0),A} X^b_{(0),B}$ consists of three basic terms:

$$\gamma_{AB} = X_{(0),A} \cdot X_{(0),B} + 2X_{(0),(A} \cdot \delta X_{,B)} + h_{ab} X^a_{(0),A} X^b_{(0),B}, \quad (\text{D.1})$$

with $X^a_{(0)}$ and δX^a as given in (C.1) and (C.3). In particular, $X_{(0),u} \cdot X_{(0),u} = X_{(0),u} \cdot X_{(0),u} = 0$ and $X_{(0),u} \cdot X_{(0),v} = \left(\frac{1}{4\pi} L_0\right)^2$, while

$$2X_{(0),u} \cdot \delta X_{,u} = 2\mu \left(\frac{1}{4\pi} L_0\right)^2 [\kappa(u - \frac{1}{2}\pi) - (F_u + A_u)], \quad (\text{D.2})$$

$$2X_{(0),(u} \cdot \delta X_{,v)} = \mu \left(\frac{1}{4\pi} L_0\right)^2 [\kappa(u - \frac{1}{2}\pi) - 2\kappa(u + v) - F_u - (F_v + A_v)] \quad (\text{D.3})$$

and

$$2X_{(0),v} \cdot \delta X_{,v} = -2\mu \left(\frac{1}{4\pi} L_0\right)^2 (G + F_v) \quad (\text{D.4})$$

for $u > 0$.

One of the components of the term proportional to h_{ab} , namely

$$h_{ab}X_{(0),u}^a X_{(0),u}^b = 16\mu(\frac{1}{4\pi}L_0)^2[\ln(W + 2u - 2\pi) - \ln(2u)], \quad (\text{D.5})$$

has already been evaluated in Appendix C. The other two components are:

$$h_{ab}X_{(0),u}^a X_{(0),v}^b = -8\mu(\frac{1}{4\pi}L_0)^2\{K(u) \cos 2u + \Sigma(u) \sin 2u - [\ln(W + 2u - 2\pi) - \ln(2u)]\} \quad (\text{D.6})$$

and

$$\begin{aligned} h_{ab}X_{(0),v}^a X_{(0),v}^b &= -8\mu(\frac{1}{4\pi}L_0)^2\{\gamma_E - \text{Ci } W + K(u) \cos 2u + \Sigma(u) \sin 2u + \ln W \\ &\quad - [\ln(W + 2u - 2\pi) - \ln(2u)]\}. \end{aligned} \quad (\text{D.7})$$

So

$$\gamma_{uu} = (\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)} [-2\mu\{F_u + A_u - \kappa(u - \frac{1}{2}\pi)\} + 16\mu\{\ln(W + 2u - 2\pi) - \ln(2u)\}], \quad (\text{D.8})$$

$$\begin{aligned} \gamma_{uv} &= (\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)} [1 - \mu\{F_u + F_v + A_v - \kappa(u - \frac{1}{2}\pi)\} \\ &\quad - 8\mu\{-\ln(W + 2u - 2\pi) + \ln(2u) + K(u) \cos 2u + \Sigma(u) \sin 2u\}] \end{aligned} \quad (\text{D.9})$$

and

$$\begin{aligned} \gamma_{vv} &= (\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)} [-2\mu(G + F_v) \\ &\quad - 8\mu\{\gamma_E - \text{Ci } W + \ln W - \ln(W + 2u - 2\pi) + \ln(2u) + K(u) \cos 2u + \Sigma(u) \sin 2u\}] \end{aligned} \quad (\text{D.10})$$

to linear order in μ , provided that $\kappa\mu|u + v| \ll 1$. Here, the factor of $e^{-2\kappa\mu(u+v)}$ inserted into γ_{uv} absorbs the term $-2\kappa(u + v)$ in (D.3), while multiplying γ_{uu} and γ_{vv} by $e^{-2\kappa\mu(u+v)}$ commits an error of order μ^2 .

Since W is a function of u alone by virtue of (6.9), the requirement that γ_{AB} take on the self-similar form $(\frac{1}{4\pi}L_0)^2 e^{-2\kappa\mu(u+v)} \bar{\gamma}_{AB}(u)$ to linear order in μ , as explained in Section 4, places three constraints on the correction functions F , G and A , namely:

$$F_{uv} + A_{uv} = 0, \quad F_{uv} + F_{vv} + A_{vv} = 0 \quad \text{and} \quad G_v + F_{vv} = 0. \quad (\text{D.11})$$

It should be noted that although equations (D.8)-(D.10) were derived on the assumption that $\kappa\mu|u + v| \ll 1$, they do in fact hold for all values of $u + v$.

This can be seen by first taking the corrected position vector (4.20) and using this to calculate $X_{,A} \cdot X_{,B}$, with the multiplicative scale factor $e^{-2\kappa\mu(u+v)}$ retained but all other terms truncated at order μ . This recovers the contribution of $X_{(0),A} \cdot X_{(0),B} + 2X_{(0),A} \cdot \delta X_{,B}$ to (D.8)-(D.10) as shown.

The contribution of the remaining term $h_{ab}X_{(0),A}^a X_{(0),B}^b$ to γ_{AB} – where $X_{(0),A}^a$ is given by (5.1) – can be evaluated by referring back to the discussion of rotating self-similarity in Section 5.1. As was demonstrated there, the value of h_{ab} at an arbitrary point

$$x^a = X_{(0)}^a(u, v) \equiv \frac{1}{4\pi} L_0 [t_L (1 - e^{-\kappa\mu(u+v)}), e^{-\kappa\mu(u+v)} \mathbf{r}(u, v)]^a \quad (\text{D.12})$$

on the worldsheet is related to its value $\bar{h}_{ab}$ at the image point

$$\bar{x}^a = \frac{1}{4\pi} L_0 e^{\kappa\mu\psi - \kappa\mu(u+v)} [|\mathbf{r}(u, v)|, R(-\psi)\mathbf{r}(u, v)]^a \quad (\text{D.13})$$

on the future light cone F_0 of the origin through the formula

$$h_{ab} = R_a^c(\psi) \bar{h}_{cd} R_b^d(\psi), \quad (\text{D.14})$$

where, from (5.13),

$$\psi = u + v - (\kappa\mu)^{-1} \ln(1 + \kappa\mu|\mathbf{r}(u, v)|). \quad (\text{D.15})$$

Here, the image point

$$\bar{x}^a = \frac{1}{4\pi} L_0 (1 + \kappa\mu|\mathbf{r}(u, v)|)^{-1} [|\mathbf{r}(u, v)|, R(-\psi)\mathbf{r}(u, v)]^a \quad (\text{D.16})$$

is just the point $X_{(0)}^a(\bar{u}, \bar{v})$ on the worldsheet with $\bar{u} = u$ and $\bar{v} = v - \psi \equiv (\kappa\mu)^{-1} \ln(1 + \kappa\mu|\mathbf{r}(u, v)|) - u$, and it was shown in Section 5.2 that $\kappa\mu|\bar{u} + \bar{v}| \ll 1$ for reasonable values of μ . So

$$X_{(0),A}^a(u, v) = X_{(0),A}^a(\bar{u}, \bar{v} + \psi) = e^{-\kappa\mu(u+v)} R_b^a(\psi) X_{(0),A}^b(\bar{u}, \bar{v}) \quad (\text{D.17})$$

to leading order in μ , and therefore

$$h_{ab} X_{(0),A}^a X_{(0),B}^b |_{(u,v)} = e^{-2\kappa\mu(u+v)} \bar{h}_{ab} X_{(0),A}^a X_{(0),B}^b |_{(\bar{u}, \bar{v})}. \quad (\text{D.18})$$

That is, to linear order in μ the term $h_{ab} X_{(0),A}^a X_{(0),B}^b$ at a general point on the worldsheet differs from its value at the image point $\bar{x}^a$ only by a multiplicative factor of $e^{-2\kappa\mu(u+v)}$, and equations (D.8)-(D.10) are recovered in full.

D.2. The length of the loop as measured along curves of constant scale factor

Equations (D.8)-(D.10) can be used to calculate the length of the evaporating loop as measured along any curve of constant scale factor $e^{-\kappa\mu(u+v)}$. The length L_{csf} of the loop along the curve $u + v = u_* + v_*$ is given by

$$L_{\text{csf}} = 2 \int_0^\pi (2\gamma_{uv} - \gamma_{uu} - \gamma_{vv})^{1/2} |_{v=u_*+v_*-u} du, \quad (\text{D.19})$$

where, to linear order in μ ,

$$\begin{aligned} (2\gamma_{uv} - \gamma_{uu} - \gamma_{vv})^{1/2} &= \sqrt{2}(\frac{1}{4\pi}L_0)e^{-\kappa\mu(u+v)}[1 + \frac{1}{2}\mu(G + A_u) + 2\mu\{\gamma_E - \text{Ci } W \\ &\quad + \ln(2uW) - \ln(W + 2u - 2\pi) - K(u) \cos 2u - \Sigma(u) \sin 2u\}]. \end{aligned} \quad (\text{D.20})$$

Hence, in view of the expression (6.50) for G :

$$\begin{aligned} L_{\text{csf}} &= 2\sqrt{2}(\frac{1}{4\pi}L_0)e^{-\kappa\mu(u_*+v_*)} \int_0^\pi [1 + \frac{1}{2}\mu(\kappa u + C_1 + A_u) \\ &\quad + 2\mu\{\gamma_E + \ln W - \text{Ci } W - 2K(u) \cos 2u - 2\Sigma(u) \sin 2u\}] du, \end{aligned} \quad (\text{D.21})$$

where

$$\begin{aligned} &\int [\ln W - \text{Ci } W - 2K(u) \cos 2u - 2\Sigma(u) \sin 2u] du \\ &= W^{-1}(2 + \ln W)(1 - \cos W) - \frac{1}{2}W(1 + \ln W) - \frac{1}{2}W^{-1}(1 - \cos 2W) - \frac{3}{2}\sin W \\ &\quad - (u - \pi) \text{Ci } W - \text{Si } W + \text{Si}(2W) - [K(u) \sin 2u - \Sigma(u) \cos 2u] \end{aligned} \quad (\text{D.22})$$

and so

$$\int_0^\pi [\ln W - \text{Ci } W - 2K(u) \cos 2u - 2\Sigma(u) \sin 2u] du = \pi \ln 2\pi + \pi - \pi \text{Ci}(2\pi) - \text{Si}(4\pi). \quad (\text{D.23})$$

Furthermore, it follows from (6.60) and (6.55) that

$$\int_0^\pi \kappa u du = \frac{1}{2}\mu\kappa\pi^2 = 2\pi \ln 2\pi + 2\pi\gamma_E - 2\pi \text{Ci}(2\pi) \quad (\text{D.24})$$

and

$$\int_0^\pi A_u du = A(\pi) - A(0) = 24\pi - 16\pi \ln 2\pi - 4 \operatorname{Si}(2\pi). \quad (\text{D.25})$$

Substituting (D.23), (D.24) and (D.25) into (D.21) therefore gives:

$$L_{\text{csf}} = \left(\frac{1}{\sqrt{2}}L_0\right)e^{-\kappa\mu(u_*+v_*)}\left[1+\frac{1}{2}\mu\{C_1+28+6\gamma_E-10\ln 2\pi-6\operatorname{Ci}(2\pi)-4\pi^{-1}\operatorname{Si}(2\pi)-4\pi^{-1}\operatorname{Si}(4\pi)\}\right]. \quad (\text{D.26})$$

Note here that $L_{\text{csf}} \rightarrow \frac{1}{\sqrt{2}}L_0$ in the flat-space limit $\mu \rightarrow 0$. This is to be expected, as the curves of constant scale factor are curves of constant t in this limit, and the speed $|\partial_t \mathbf{X}|$ is equal to $1/\sqrt{2}$ at all points on the unperturbed ACO loop (except at the kink points at $u = 0$ and π , where it is undefined). The measured length $L_{\text{csf}} = \frac{1}{\sqrt{2}}L_0$ is therefore just the Lorentz-contracted fundamental length L_0 .

If μ is non-zero then L_{csf} will deviate from its ACO value $\frac{1}{\sqrt{2}}L_0$ on the curve $u + v = 0$ by a term of order μ unless C_1 takes on the particular value

$$C_1 = -28 - 6\gamma_E + 10\ln 2\pi + 6\operatorname{Ci}(2\pi) + 4\pi^{-1}\operatorname{Si}(2\pi) + 4\pi^{-1}\operatorname{Si}(4\pi) \approx -9.5143. \quad (\text{D.27})$$

There is of course no necessary reason to choose this value for C_1 , but it bears repeating that different values of C_1 will lead to different “dressed” values of L_{csf} along any given curve of constant $u + v$.

D.3. Invariant length of the evaporating loop

Another measure of the length of the loop that is potentially of interest here is the *invariant length* L_I , which is a gauge-invariant quantity defined at each point (u_*, v_*) on the worldsheet by

$$L_I(u_*, v_*) = 2 \left(\int_{\mathcal{D}(u_*, v_*)} \gamma^{1/2} du dv \right)^{1/2}, \quad (\text{D.28})$$

where $\mathcal{D}(u_*, v_*)$ is the closure of the subset of the worldsheet that is causally disconnected from the reference point (u_*, v_*) [31]. Calculating L_I to order μ at a general point on the trajectory (4.20) of the evaporating ACO loop is a lengthy process whose details are of marginal importance, so I will offer only an abbreviated summary here.

In the flat-space limit $\mu = 0$ the domain $\mathcal{D}(u_*, v_*)$ is bounded by the null curves $u = u_*$ and $v = v_*$ which, in view of the identification of (u, v) with $(u +$

$2\pi k, v - 2\pi k$) for all integers k , intersect at the points $(u_*, v_* + 2\pi) \equiv (u_* + 2\pi, v_*)$ and $(u_*, v_* - 2\pi) \equiv (u_* - 2\pi, v_*)$. The boundary of $\mathcal{D}(u_*, v_*)$ is therefore completed by the curves

$$v = u + 2\pi - u_* + v_* \quad \text{and} \quad v = u - 2\pi - u_* + v_* \quad (\text{D.29})$$

and so

$$\begin{aligned} \lim_{\mu \rightarrow 0} L_I(u_*, v_*) &= 2 \left(\int_{u_* - 2\pi}^{u_*} \int_{v_*}^{u + 2\pi - u_* + v_*} \left(\frac{1}{4\pi} L_0\right)^2 dv du + \int_{u_*}^{u_* + 2\pi} \int_{u - 2\pi - u_* + v_*}^{v_*} \left(\frac{1}{4\pi} L_0\right)^2 dv du \right)^{1/2} \\ &= L_0 \end{aligned} \quad (\text{D.30})$$

In order to calculate L_I when μ is non-zero, it is useful to first transform to a set of gauge coordinates (u', v') with the property that the curves $u' = u_*$ and $v' = v_*$ are (to order μ) null curves through the reference point $(u, v) = (u_*, v_*)$. A suitable choice is

$$u' = u - \mu(v - v_*)\mathcal{G}(u) \quad \text{and} \quad v' = v - \mu\mathcal{F}(u; u_*), \quad (\text{D.31})$$

where

$$\begin{aligned} \mathcal{G}(u) &\equiv G(u) + 4\{\gamma_E - \text{Ci } W + \ln W - \ln(W + 2u - 2\pi) + \ln(2u) \\ &\quad + K(u) \cos 2u + \Sigma(u) \sin 2u\} \\ &= \kappa u + 4(\gamma_E - \text{Ci } W + \ln W) + C_1 \end{aligned} \quad (\text{D.32})$$

and

$$\mathcal{F}(u; u_*) \equiv \int_{u_*}^u [\{F_u + A_u - \kappa(u - \tfrac{1}{2}\pi)\} - 8\{\ln(W + 2u - 2\pi) - \ln(2u)\}] du. \quad (\text{D.33})$$

In the primed gauge the 2-metric components (D.8)-(D.10) become $\gamma_{u'u'} = \gamma_{v'v'} = 0$ and

$$\begin{aligned} \gamma^{1/2} &\equiv \gamma_{u'v'} = \left(\frac{1}{4\pi} L_0\right)^2 e^{-2\kappa\mu(u+v)} [1 - \mu\{F_u - \kappa(u - \tfrac{1}{2}\pi)\} + \mu(v - v_*)\partial_u \mathcal{G}(u) \\ &\quad - 8\mu\{-\ln(W + 2u - 2\pi) + \ln(2u) + K(u) \cos 2u + \Sigma(u) \sin 2u\}] \end{aligned} \quad (\text{D.34})$$

to linear order in μ .¹⁴

¹⁴Strictly speaking, the functions of u on the right-hand side of (D.32), (D.33) and (D.34) should be the half-range periodic extensions of the expressions shown. However, all three equations are accurate if u and u_* both lie in $[0, \pi]$.

Recalling again that the point (u, v) is identified with $(u \pm 2\pi, v \mp 2\pi)$, the curves $u' = u_*$ and $v' = v_*$ intersect at the points

$$(u, v) = (u_* - \Delta u, v_* + 2\pi - \Delta v) \equiv (u_* + 2\pi - \Delta u, v_* - \Delta v) \quad (\text{D.35})$$

and

$$(u, v) = (u_* + \Delta u, v_* - 2\pi + \Delta v) \equiv (u_* - 2\pi + \Delta u, v_* + \Delta v), \quad (\text{D.36})$$

where

$$\Delta u \equiv -2\pi\mu\mathcal{G}(u_*) \quad (\text{D.37})$$

and

$$\begin{aligned} \Delta v &\equiv -\mu\mathcal{F}(u_* + 2\pi; u_*) \\ &= -2\mu \int_0^\pi [\{F_u + A_u - \kappa(u - \tfrac{1}{2}\pi)\} - 8\{\ln(W + 2u - 2\pi) - \ln(2u)\}] du \\ &= -[16\pi - 8\text{Si}(2\pi)]\mu \approx -38.92\mu. \end{aligned} \quad (\text{D.38})$$

The deviation of v by the amount $\pm 38.92\mu$ from its flat-space values v_* and $v_* \pm 2\pi$ at the four intersection points accounts for the rotational phase shift of the same magnitude, mentioned at the end of Section 3, that is induced by gravitational back reaction on the ACO loop over the course of a single oscillation period.

The boundary of $\mathcal{D}(u_*, v_*)$ is completed by the straight line segments (in (u, v) coordinates) joining $(u_* - \Delta u, v_* + 2\pi - \Delta v)$ to $(u_* - 2\pi + \Delta u, v_* + \Delta v)$, and $(u_* + 2\pi - \Delta u, v_* - \Delta v)$ to $(u_* + \Delta u, v_* - 2\pi + \Delta v)$. In the primed gauge, these segments become

$$v' = v_* + (u' - u_* + 2\pi)[1 - \pi^{-1}\Delta v + \pi^{-1}\Delta u + \mu\mathcal{G}(u')] + \Delta v - \Delta u - \mu\mathcal{F}(u'; u_*) \quad (\text{D.39})$$

and

$$v' = v_* + (u' - u_* - 2\pi)[1 - \pi^{-1}\Delta v + \pi^{-1}\Delta u + \mu\mathcal{G}(u')] - \Delta v + \Delta u - \mu\mathcal{F}(u'; u_*) \quad (\text{D.40})$$

to first order in μ .

The equation (D.28) therefore becomes:

$$L_I(u_*, v_*) = 2 \left(\int_{u_* - 2\pi + \Delta u}^{u_*} \int_{v_*}^{v_1(u')} \gamma^{1/2} dv' du' + \int_{u_*}^{u_* + 2\pi - \Delta u} \int_{v_2(u')}^{v_*} \gamma^{1/2} dv' du' \right)^{1/2} \quad (\text{D.41})$$

where v_1 and v_2 are the functions on the right-hand sides of (D.39) and (D.40) respectively, and the form of $\gamma^{1/2}$ is given in (D.34), with u and v everywhere replaced by u' and v' . If the scale factor $e^{-2\mu(u+v)}$ in (D.34) is expanded as $e^{-2\mu(u_*+v_*)}[1 - 2\mu(u' - u_* + v' - v_*)]$ then the integrals inside the square root in (D.41) can be broken into two parts:

$$\left(\frac{1}{4\pi}L_0\right)^2 e^{-2\kappa\mu(u_*+v_*)} \left(\int_{u_*-2\pi+\Delta u}^{u_*} [v_1(u') - v_*] du' + \int_{u_*}^{u_*+2\pi-\Delta u} [v_* - v_2(u')] du' \right) \quad (\text{D.42})$$

and

$$\left(\frac{1}{4\pi}L_0\right)^2 e^{-2\kappa\mu(u_*+v_*)} \left(\int_{u_*-2\pi}^{u_*} \int_{v_*}^{u'+2\pi-u_*+v_*} \mu \Delta dv' du' + \int_{u_*}^{u_*+2\pi} \int_{u'-2\pi-u_*+v_*}^{v_*} \mu \Delta dv' du' \right), \quad (\text{D.43})$$

where

$$\begin{aligned} \Delta \equiv & -2(u' - u_* + v' - v_*) - \{F_{u'} - \kappa(u' - \tfrac{1}{2}\pi)\} + (v' - v_*)\partial_{u'}\mathcal{G}(u') \\ & -8\{-\ln(W + 2u' - 2\pi) + \ln(2u') + K(u')\cos 2u' + \Sigma(u')\sin 2u'\}. \end{aligned} \quad (\text{D.44})$$

Now, it is easily seen that the contribution of the term $-2(u' - u_* + v' - v_*)$ in Δ to the integrals in (D.43) is zero, and since the remaining terms in Δ are all half-range periodic functions of u' it turns out that

$$\begin{aligned} & \int_{u_*-2\pi}^{u_*} \int_{v_*}^{u'+2\pi-u_*+v_*} \mu \Delta dv' du' + \int_{u_*}^{u_*+2\pi} \int_{u'-2\pi-u_*+v_*}^{v_*} \mu \Delta dv' du' \\ = & 2\pi\mu \int_{u_*}^{u_*+2\pi} (u' - u_* - \pi)\partial_{u'}\mathcal{G}(u') du - 4\pi\mu \int_0^\pi \{F_u - \kappa(u - \tfrac{1}{2}\pi)\} du \\ & -32\pi\mu \int_0^\pi \{-\ln(W + 2u' - 2\pi) + \ln(2u') + K(u')\cos 2u' + \Sigma(u')\sin 2u'\} du'. \end{aligned} \quad (\text{D.45})$$

Furthermore, the contribution of the second integral here is zero, as $F(\pi) = F(0)$ and $u - \frac{1}{2}\pi$ integrates to zero.

Turning to the integrals in (D.42), the terms in $v_1(u') - v_*$ and $v_* - v_2(u')$ of order μ^0 contribute $4\pi^2$ to the integrals as expected, while the terms proportional to Δu and Δv integrate to give zero to order μ . Also, since $\mathcal{G}$ is half-range periodic

in u' , and $\mu\mathcal{F}(u'; u_*) - \mu\mathcal{F}(u' - 2\pi; u_*) = -\Delta v$ from (D.33) and (D.38), it turns out that

$$\int_{u_*-2\pi+\Delta u}^{u_*} [v_1(u') - v_*] du' + \int_{u_*}^{u_*+2\pi-\Delta u} [v_* - v_2(u')] du' = 4\pi^2 + 2\pi\mu \int_{u_*}^{u_*+2\pi} \mathcal{G}(u') du' - 2\pi\Delta v. \quad (\text{D.46})$$

Combining (D.45) and (D.46) gives:

$$\begin{aligned} & \int_{u_*-2\pi+\Delta u}^{u_*} \int_{v_*}^{v_1(u')} \gamma^{1/2} dv' du' + \int_{u_*}^{u_*+2\pi-\Delta u} \int_{v_2(u')}^{v_*} \gamma^{1/2} dv' du' \\ &= 4\pi^2 + 4\pi^2\mu\mathcal{G}(u_*) + 2\pi\mu[16\pi - 8\text{Si}(2\pi)] \\ & \quad - 32\pi\mu \int_0^\pi \{-\ln(W + 2u' - 2\pi) + \ln(2u') + K(u') \cos 2u' + \Sigma(u') \sin 2u'\} du' \end{aligned} \quad (\text{D.47})$$

where the value of the integral in the final term is $2\pi \ln 2\pi + \text{Si}(2\pi) - 3\pi$.

So, to first order in μ , the invariant length of the loop is:

$$\begin{aligned} L_I(u_*, v_*) &= 2\left(\frac{1}{4\pi}L_0\right)e^{-\kappa\mu(u_*+v_*)}\{4\pi^2 + 4\pi^2\mu[\kappa u_* + 4(\gamma_E - \text{Ci } W_* + \ln W_*) + C_1] \\ & \quad - 16\pi\mu[-8\pi + 4\pi \ln 2\pi + 3\text{Si}(2\pi)]\}^{1/2} \\ &= L_0 e^{-\kappa\mu(u_*+v_*)}\{1 + \mu[\frac{1}{2}\kappa u_* + 2(\gamma_E - \text{Ci } W_* + \ln W_*) + C_1] \\ & \quad - 2\mu[-8 + 4\ln 2\pi + 3\pi^{-1}\text{Si}(2\pi)]\} \end{aligned} \quad (\text{D.48})$$

for $u_* \in (0, \pi)$. For values of u_* outside this range, the term multiplying the scale factor $e^{-\kappa\mu(u_*+v_*)}$ should be replaced by its half-range periodic extension. Furthermore the u_* -dependent part of this term, $\frac{1}{2}\kappa u_* + 2(\gamma_E - \text{Ci } W_* + \ln W_*)$, tends to $2(\gamma_E - \text{Ci } 2\pi + \ln 2\pi)$ as $u_* \rightarrow 0^+$, and to $\frac{1}{2}\kappa\pi$ as $u_* \rightarrow \pi^-$. In view of the value (6.60) deduced for κ in Section 6.3, the two limits are the same, and so L_I is continuous for all values of u_* and v_* .

References

- [1] Vilenkin A and Shellard E P S 1990 *Cosmic strings and other topological defects* (Cambridge: Cambridge University Press)
- [2] Geroch R and Traschen J 1987 *Phys. Rev. D* **36** 1017
- [3] Gott J R 1985 *Astrophys. J.* **288** 422

- [4] Hiscock W A 1985 *Phys. Rev. D* **31** 3288
- [5] Xanthopoulos B 1986 *Phys. Lett.* **178B** 163
- [6] Garriga J and Verdaguer E 1987 *Phys. Rev. D* **36** 2250
- [7] Garfinkle D 1990 *Phys. Rev. D* **41** 1112
- [8] Frolov V P and Garfinkle D 1990 *Phys. Rev. D* **42** 3980
- [9] Hawking S W 1990 *Phys. Lett.* **246B** 36
- [10] Frolov V P, Israel W and Unruh W G 1989 *Phys. Rev. D* **39** 1084
- [11] Quashnock J M and Spergel D N 1990 *Phys. Rev. D* **42** 2505
- [12] Anderson M R 2005 *Class. Quant. Grav.* **22** 2539
- [13] Allen B, Casper P and Ottewill A 1994 *Phys. Rev. D* **50** 3703
- [14] Nambu Y 1971 *Phys. Rev. D* **4** 1193
- [15] Goto T 1971 *Prog. Theor. Phys.* **46** 1560
- [16] Battye R A and Carter B 1995 *Phys. Lett. B* **357** 29
- [17] Carter B and Battye R A 1998 *Phys. Lett. B* **430** 49
- [18] Allen B and Shellard E P S 1992 *Phys. Rev. D* **45** 1898
- [19] Allen B and Casper P 1994 *Phys. Rev. D* **50** 2496
- [20] Barrabès C and Israel W 1991 *Phys. Rev. D* **43** 1129
- [21] Poisson E 2002 arXiv:gr-qc/0207101v1
- [22] Kibble T W B and Turok N 1982 *Phys. Lett. B* **116** 141
- [23] Anderson M R 2003 *The Mathematical Theory of Cosmic Strings* (Bristol: IOP Publishing)
- [24] Vanchurin V, Olum K D and Vilenkin A 2006 *Phys. Rev. D* **74** 063527
- [25] Ringeval C, Sakellariadou M and Bouchet F R 2007 *JCAP* 0702:023

- [26] Vanchurin V 2008 *Phys. Rev. D* **77** 063532
- [27] Dubath F, Polchinski J and Rocha J V 2008 *Phys. Rev. D* **77** 123528
- [28] Vincent G R, Hindmarsh M and Sakellariadou M 1997 *Phys. Rev. D* **56** 637
- [29] Martins C J A P and Shellard E P S 2006 *Phys. Rev. D* **73** 043515
- [30] Polchinski J and Rocha J V 2007 *Phys. Rev. D* **75** 123503
- [31] Anderson M R 1990 *Phys. Rev. D* **41** 3612

LIST OF FIGURE CAPTIONS:

Figure 1: y - z projection of the Allen-Casper-Ottewill loop.

Figure 2: Schematic representation of the evaporation of the ACO loop.

Figure 3: Schematic picture of the relationship between the field point $[t, \mathbf{x}]$ and its image point $[\bar{t}, \bar{\mathbf{x}}]$.

Figure 4a: Variation of $h_{tt}/(4\mu)$ with Z along the $\bar{z}$ -axis.

Figure 4b: Variation of $h_{tx}/(4\mu)$ with Z along the $\bar{z}$ -axis.

Figure 4c: Variation of $h_{ty}/(4\mu)$ with Z along the $\bar{z}$ -axis.

Figure 4d: Variation of $h_{zx}/(4\mu)$ with Z along the $\bar{z}$ -axis.

Figure 4e: Variation of $h_{zy}/(4\mu)$ with Z along the $\bar{z}$ -axis.

Figure 5a: The correction G as a function of u if $C_1 \approx -9.5143$.

Figure 5b: The combination $F + \text{sgn}(u) A$ as a function of u if $C_2 \approx 15.403$.

Figure 5c: The “natural” correction function $A(u) = -\frac{2\lambda}{\pi}(|u| - \frac{1}{2}\pi)$.

Figure 5d: The correction F as a function of u if A has its natural form.

Figure 6: The Kibble-Turok sphere of the ACO loop and two possible types of perturbation.




Figure 1




Figure 2




Figure 3




Figure 4a




Figure 4b




Figure 4c




Figure 4d




Figure 4e




Figure 5a




Figure 5b




Figure 5c




Figure 5d




Figure 6